



King's Research Portal

DOI:

[10.1001/jamapsychiatry.2019.2859](https://doi.org/10.1001/jamapsychiatry.2019.2859)

[10.1001/jamapsychiatry.2019.2859](https://doi.org/10.1001/jamapsychiatry.2019.2859)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Dragioti, E., Solmi, M., Favaro, A., Fusar-Poli, P., Dazzan, P., Thompson, T., Stubbs, B., Firth, J., Fornaro, M., Tsartalis, D., Carvalho, A. F., Vieta, E., McGuire, P., Young, A. H., Shin, J. I., Correll, C. U., & Evangelou, E. (2019). Association of Antidepressant Use With Adverse Health Outcomes: A Systematic Umbrella Review. *JAMA Psychiatry*, 76(12), 1241-1255. <https://doi.org/10.1001/jamapsychiatry.2019.2859>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Title sheet

Title: Adverse health outcomes in antidepressants users.

Subtitle: An umbrella review of 45 meta-analyses of observational studies.

Elena Dragioti*, PhD, ^{1,2} Marco Solmi*, MD, PhD, ^{3,4,5} Angela Favaro^{3,4}, MD, PhD, Paolo Fusar-Poli^{5,6,7}, MD, PhD, Paola Dazzan⁵, MD, PhD, Trevor Thompson⁸, PhD, Brendon Stubbs^{9,10}, PhD, Joseph Firth^{11,12,13}, PhD, Michele Fornaro¹⁴, MD, PhD, Dimitrios Tsartalis¹⁵, MD, PhD, Andre F. Carvalho¹⁶, MD, PhD, Eduard Vieta¹⁷, MD, PhD, Philip McGuire⁵, MD, PhD, Allan H. Young^{10,18}, FRCPsych, Jae Il Shin¹⁹, MD, PhD, Christoph U. Correll, MD^{20,21,22,23}, Evangelos Evangelou^{24,25}, PhD

¹ Pain and Rehabilitation center and Department of Medicine and Health Sciences (IMH), Faculty of Health Sciences University of Linköping, SE- 581 85 Linköping, Sweden

² Department of Hygiene and Epidemiology, School of Medicine, University of Ioannina, University Campus, 45110 Ioannina, Greece

³Neurosciences Department, University of Padua, Padua, Italy

⁴Neuroscience Center, University of Padua, Padua, Italy

⁵Early Psychosis: Interventions and Clinical-detection (EPIC) lab, Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom;

⁶ OASIS service, South London and Maudsley NHS Foundation Trust, London, United Kingdom;

⁷ Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy;

⁸ Department of Psychology, Social Work and Counselling, University of Greenwich, UK

⁹ Physiotherapy Department, South London and Maudsley NHS Foundation Trust, London, United Kingdom.

¹⁰ Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom.

¹¹ NICM Health Research Institute, School of Science and Health, University of Western Sydney, Sydney, Australia.

¹² Division of Psychology and Mental Health, Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK.

¹³ Centre for Youth Mental Health, University of Melbourne, Melbourne, Australia.

¹⁴ Department of Neuroscience, Reproductive Sciences and Dentistry, Federico II University, Naples, Italy

¹⁵ Department of Clinical Physiology, Linköping University, Linköping, SE-58183, Linköping, Sweden.

¹⁶ Centre for Addiction and Mental Health (CAMH) and Department of Psychiatry, University of Toronto, Toronto, ON, Canada.

¹⁷ Department of Psychiatry and Psychology, Hospital Clinic, Institute of Neuroscience, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Catalonia, Spain.

¹⁸ South London and Maudsley NHS Foundation Trust, Bethlem Royal Hospital, Monks Orchard Road, Beckenham, Kent, BR3 3BX, United Kingdom

¹⁹ Department of Pediatrics, Yonsei University College of Medicine, Seoul, Republic of Korea

²⁰ Department of Psychiatry, Zucker Hillside Hospital, Glen Oaks, NY, USA.

²¹ Department of Psychiatry and Molecular Medicine, Hofstra Northwell School of Medicine, Hempstead, NY, USA.

²² Center for Psychiatric Neuroscience, Feinstein Institute for Medical Research, Manhasset, NY, USA.

²³ Campus Virchow-Klinikum, Charité-Universitätsmedizin Berlin, and Department of Child and Adolescent Psychiatry, Berlin Institute of Health, Berlin, Germany.

²⁴ Department of Epidemiology and Biostatistics, Imperial College London, London, UK.

²⁵ Department of Hygiene and Epidemiology, University of Ioannina Medical School, Ioannina, Greece.

Correspondence to: Elena Dragioti, PhD, Senior Lecturer, Pain and Rehabilitation center and Department of Medicine and Health Sciences (IMH), Faculty of Health Sciences University of Linköping, SE- 581 85 Linköping, Sweden and Department of Hygiene and Epidemiology, School of Medicine, University of Ioannina, University Campus, 45110 Ioannina, Greece, E-mail: elena.dragioti@liu.se

* Both authors contributed equally to the manuscript as joint first authors

Words: 3533; References:87; Tables: 3; Figure 2

Date of revision: June 20th 2019

Abstract

Importance: Antidepressant use is increasing worldwide. Contrasting evidence from meta-analyses is available on their safety.

Objective: To grade the evidence across meta-analyses of observational studies assessing the association between antidepressants and adverse health outcomes, using an umbrella review approach.

Data Sources: We searched PubMed, Scopus, and PsycINFO, up to April 5, 2019.

Evidence Review: Only meta-analyses of observational studies with a cohort or case-control study design were eligible. Two independent reviewers recorded the data and assessed the methodological quality of the selected meta-analyses. Evidence was ranked as convincing, highly suggestive, suggestive, weak, and not significant according to established criteria.

Results: Forty-five eligible meta-analyses describing 120 associations including data from 1012 individual effect size estimates were included, after scrutinizing 252 potential articles. Seventy-four of the 120 associations (61.7%) had a nominally statistically significant effect ($p \leq 0.05$) using random-effects models. Large heterogeneity ($I^2 > 50\%$) was present (43.3%), while small-study effects (14.2%) and excess significance bias (7.5%) were less common. While convincing meta-analytic evidence emerged from both main and sensitivity analysis for the association between antidepressants and risk of suicide attempt/completion in children/adolescents (9.4%, OR=1.92, 95%CI=1.51-2.44 in adjusted studies, SSRIs only, and in high-quality studies, 3.6% OR=1.88, 95%CI=1.47-2.40), autism spectrum disorders (ASD) with antidepressants before (0.8%, RR=1.48, 95%CI=1.29-1.71 in any antidepressant, adjusted studies, and high-quality studies) and during pregnancy (0.9%, OR=1.84, 95%CI=1.60-2.11 in both high-quality studies and SSRIs only, and 0.1% OR=1.80, 95%CI=1.54-2.10 in European studies), pre-term birth in prospective cohort studies (0.4%, RR=1.87, 95%CI=1.52-2.30), and in studies of mixed antidepressants (0.4%, RR=1.59, 95%CI=1.31-1.93), and low APGAR scores in SSRIs only studies (5.7%, SMD=-0.27, 95%CI -0.37

to -0.16), none of these associations remained at convincing evidence after sensitivity analysis for confounding by indication.

Conclusions and Relevance: Most putative adverse health outcomes associated with antidepressants are not supported by convincing evidence, and the few of those supported by convincing evidence are affected by confounding by indication. Antidepressants appear to be overall safe for the treatment of psychiatric disorders. More studies matching for underlying disease are needed to clarify the degree of confounding by indication and other biases. No absolute contraindication to antidepressants emerges from this umbrella review.

Registration: PROSPERO registration 2018: CRD42018103462.

Key words: antidepressants; adverse outcomes; umbrella review; meta-analysis; safety

Key Points

Question: Are antidepressants associated with adverse health outcomes and how strong is the evidence in the published meta-analyses of real-world data?

Findings: In this systematic umbrella review of 45 meta-analyses of observational studies, convincing evidence was found for the association between antidepressants and suicide behavior/completion in individuals aged <19 years old and for autism risk in the offspring. However, none of these associations remained at convincing evidence after sensitivity analysis for confounding by indication.

Meaning Claimed adverse health outcomes associated with antidepressants are not supported by strong evidence and are exaggerated by confounding by indication. No absolute contraindication to the use of antidepressants is currently supported by convincing evidence.

Introduction

Accumulating evidence suggests that there is a sharp growth of antidepressants use worldwide (up to 8-10% of American adults take at least one antidepressant - rank third among prescribed and fourth among sold medications).^{1,2} Antidepressants are indicated and used for depressive disorders, anxiety disorders, post-traumatic stress disorder, premenstrual dysphoric disorder, obsessive compulsive disorder (OCD), binge eating disorder, among others.³⁻⁵

The safety profile of antidepressants is controversial. Since the Food and Drug Administration (FDA) introduced the black box warnings associating selective serotonin reuptake inhibitors (SSRIs) with an increased risk of suicidal behavior in children and adolescents,⁶ the debate about efficacy, acceptability, and the safety profile of antidepressant medications has gradually increased.⁷⁻¹¹ While evidence from randomized controlled trials (RCTs) showing antidepressants' efficacy and acceptability has been well documented, in both meta-analyses and network meta-analyses,^{4,8,10,12,13} safety assessment is inherently biased by certain methodological weaknesses of RCTs, such as small and unrepresentative samples, rare and inconsistent reporting of adverse outcomes, and short duration of exposures.^{14,15}

Observational studies provide complementary evidence to RCTs on a number of adverse health outcomes (which may only manifest in the mid- to long-term) utilizing real-world data¹⁵ linked to antidepressants, which it is not possible in RCTs.¹⁶ In particular, observational studies can provide useful evidence on medication safety since they are more representative of overall target population, for example including patients with comorbid disorders or suicidal thoughts, which are often excluded from RCTs. Also, observational studies typically have a longer follow up duration than RCTs, providing data on mid- and long-term effects of antidepressants, such as bone status, or gastro-intestinal bleeding which may not arise from short-term use.¹⁶ Several meta-analyses of observational studies assessing antidepressant safety have been published, but no attempt of quantifying their credibility has been made to date. The latter is crucial, considering the uncertainty

surrounding observational research results.¹⁷⁻¹⁹ Umbrella reviews make it feasible to summarize the evidence from multiple meta-analyses on the same topic^{20,21} and enable a grading (ranking) of evidence into convincing, highly suggestive, suggestive, weak, or not significant based on sample size, strength of the association, and the assessment of presence of biases.²²⁻²⁴ In this study, we graded the evidence from published meta-analyses of observational studies testing the association between antidepressant use and risk of adverse health outcomes.

Methods

A protocol for this study was registered on PROSPERO 2018: CRD42018103462. We used the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) recommendations²⁵ and the Meta-analysis of Observational Studies in Epidemiology (MOOSE) guidelines²⁶ to guide this umbrella review.

Search strategy and selection criteria

We searched the PubMed, Scopus, and PsycINFO, from inception to April 5, 2019, to identify systematic reviews with meta-analysis of observational studies that examined an association between any adverse health outcome and exposure to antidepressants. We performed a search strategy using a combination of terms related to antidepressants (e.g., "antidepressants", "selective serotonin reuptake inhibitors"), terms related to adverse health outcomes (e.g., "harms," "suicide," "bleeding," "autism," etc.), and to meta-analysis with no age, sex, population, and medical condition restrictions (for details, see eBox1 in the Supplement). We also hand searched the cited references of the retrieved articles and reviews.

Two reviewers (ED, MS) independently searched titles/abstracts for eligibility, and when a consensus could not be achieved a third reviewer was consulted (EE). The full-texts of potentially eligible articles were retrieved, and the same two investigators independently scrutinized each study for eligibility. Any discrepancies were resolved by the third reviewer (EE).

We included only peer-reviewed systematic reviews with meta-analysis of observational studies with only a cohort, case-control, or nested case-control study design, measuring any association between antidepressants and any adverse health outcome in any population of any age. Whenever multiple meta-analyses on the same adverse health outcome were performed (i.e., overlapping meta-analyses with the same outcome, type of antidepressant, and clinical setting or population setting), we assessed only the one that included the largest data set, as previously described.^{22,24,27} Details of the selection between overlapping meta-analyses are described in the Supplement. For each eligible meta-analysis, we considered the main analysis for all primary and secondary reported outcomes. The concordance between selected and non-selected meta-analyses was examined in a sensitivity analysis.²⁷

We excluded: 1) meta-analyses of other study designs (e.g., RCTs, cross-sectional) or meta-analysis included both observational studies and RCTs in the same analysis, 2) published in languages other than English, 3) meta-analyses of individual patient or participant data, pooled analyses that examined a non-systematic selection of observational studies, and non-systematic reviews, 4) meta-analyses of hypericum perforatum, known as perforate St John's-wort, or tryptophan, and 5) meta-analyses that provided insufficient or inadequate data for quantitative synthesis.

Data extraction

Data extraction was performed independently by two investigators (ED, MS) and disagreements were resolved by a consensus. Adverse health outcomes related to exposure to antidepressants were extracted as defined by the original authors. For each meta-analysis, we recorded information on PMID/DOI, first author, publication year, type of antidepressant, study design, age of participants, adverse health outcomes, exposure and non-exposure, illnesses examined (e.g., depression), the number of included studies, and the total sample size.

For each primary study, we recorded information on first author, year of publication, study design (i.e., cohort or case-control), number of cases and controls in case-control studies or total population in cohort studies, reported adjusted (or unadjusted, if such exists) effect sizes (ESs), i.e.; risk ratio (RR); odds ratio (OR), hazard ratio (HR), standardized mean difference (SMD), each with their 95% confidence intervals (CIs), and study location. We also captured the number and nature of adjustments, length of follow-up, study quality score, and if the studies were controlled for a psychiatric condition, i.e., confounding by indication.^{18,19}

Data were extracted independently by two investigators (ED, MS). The methodological quality of each included meta-analysis was assessed with the Assessment of multiple systematic reviews (AMSTAR) 2 tool (available at <https://amstar.ca/Amstar-2.php>), which is a recent update of AMSTAR,²⁸ by same two investigators (ED, MS). AMSTAR 2 also accounts for the quality of studies included in the meta-analysis, beyond a mere technical methodology assessment of the included meta-analysis (for details see supplement).¹⁶

Data analysis

For each association, we extracted effect sizes of individual studies included in each meta-analysis, and re-meta-analyzed them calculating the pooled ES and the 95% confidence intervals, with random-effects models to compare homogeneously analyzed results.²⁹ We did not transform the initial ESs or modify the direction of associations presented by the original authors to compare our results with the reported results from the meta-analyses. Heterogeneity was assessed with the I^2 statistic.³⁰ Additionally, we calculated the 95% prediction intervals for the summary random ESs providing the possible range in which the ESs of future studies is expected to fall.³¹

We then tested whether smaller studies yielded larger ESs than larger ones; an indication of small-study effect bias.^{24,32-34} Small-study effect bias was indicated by both Egger's regression asymmetry test ($p \leq 0.10$) and by the random effects' summary ES being larger than the ES of the largest study in each association.^{24,32-34} Finally, we assessed the existence of excess significance bias

by evaluating whether the observed number (O) of studies with nominally statistically significant results (“positive” studies, $p \leq 0.05$) were different from the expected number (E) of studies with statistically significant results.³⁴ The E of statistically significant studies per association was calculated by summing the statistical power estimates for each component study. The power estimates of each component study depend on the plausible effect size for the tested association, which we assumed to be the effect of the largest study (i.e., the smallest standard error) per association.³⁵ Excess significance bias was set at $p \leq 0.10$. This test is designed to assess whether the published meta-analyses comprise an over-representation of false positive findings.³⁴

Assessment of the credibility of the evidence

We assessed the credibility of the evidence per association provided in meta-analyses by applying several criteria in concordance with previous published umbrella reviews.^{22,23,32,33,36,37} In brief, associations that presented nominally significant random-effects summary ESs (i.e., $p \leq 0.05$) were ranked as convincing, highly suggestive, suggestive, and weak evidence based on sample size, strength of the association, and the assessment of presence of biases (Table 1; for details see Supplement). In addition, to provide an estimate of epidemiologic implication of findings, we calculated prevalence of outcomes of interest from cohort studies only (studies with case-control designs should not be considered for prevalence estimates).

Sensitivity analysis

We performed sensitivity analyses to assess whether the credibility of the evidence varied within cohort studies (both prospective and retrospective), prospective cohort studies, studies with adjustments for multiple covariates, confounding by indication, high-quality studies, classes of antidepressants (SSRIs studies, tricyclic antidepressants [TCAs] studies, other or mixed antidepressants studies), and study location of the associations ranked as convincing and highly suggestive (i.e., class I-II) in the main analysis.

Results

We identified 4,471 articles, scrutinized 252 in full text and finally included 45 meta-analyses in this umbrella review³⁸⁻⁸¹ (Figure 1), corresponding to 695 studies, 1012 estimates, and 13 putative risks (Figure 2). References of the 207 excluded articles and their reasons are provided in eTable 1.

Descriptive characteristics of the 45 eligible meta-analyses of observational studies can be found in eTable2. All meta-analyses but one had also a control group not exposed to antidepressants, in which compared the risk of gastrointestinal (GI) bleeding between mirtazapine and SSRIs.⁴⁷ The median number of adjustments in the analyses was 7 (interquartile range [IQR]=4-11, range 0-31) and the median duration of follow-up was 4 years (IQR=2-5).

Thirty-three meta-analyses (73.4%) met the moderate-quality level according to the AMSTAR2 evaluation, and eight (17.8%) were of low quality. Finally, two (4.4%) were high-quality, while two others (4.4%) were of critically low quality. The two independent investigators reached a high level of agreement (91.0%) on the quality rating.

Description of associations

Forty-five eligible meta-analyses described 120 associations including 1012 individual study estimates of adverse health outcomes related to exposure to antidepressants (see Tables 1 to 2, and eTables 3 to 5 in the Supplement), with a median number of estimates per association of 6 (IQR =4-12). Seventy-four (61.7%) of the associations concerned maternal and pregnancy-related adverse health outcomes (Figure 2). Most associations (n=80; 66.7%) concerned SSRIs or SNRIs, nine (7.5%) TCAs, and 31 (25.7%) mixed or other antidepressants.

The median number of the total population per association was 1,056,374 (IQR=152,180 – 2,215,969). The median number of cases (adverse health outcomes) per association was 12,097 (IQR=2,585–56,272) and the number of cases was >1,000 for 87 associations (72.5%).

Summary of associations

A summary of all 120 associations is presented in Tables 2 to 3 and eTables 3 to 5 in the Supplement. Seventy-four of the 120 examined associations (61.7%) had a nominally statistically

significant effect ($p \leq 0.05$) using random-effects models and only 22 of those (18.3%) reached $p \leq 1 \times 10^{-6}$. Almost all statistically significant associations indicated an increased risk for antidepressants and adverse health outcomes, except for two associations showing a protective effect of SSRI against suicide attempt/completion in adults and older adults, respectively.⁸⁰

Fifty-two associations (43.3%) had large heterogeneity ($I^2 > 50\%$), and only for 24 associations (20.0%) the 95% prediction intervals excluded the null value. In 63 associations (52.5%), the ESs of the largest study presented a nominally statistically significant effect ($p \leq 0.05$). Finally, small-study effects were found for 17 associations (14.2%), and excess significance bias was observed for nine associations (7.5%).

Grading of level of evidence of associations

Main analysis

Convincing evidence

Among the 120 associations, three (2.5%) had convincing evidence; the association between SSRIs and increased risk of suicide attempt/completion in children and adolescents,⁸⁰ and the association between any antidepressant before pregnancy and SSRIs during pregnancy and autism spectrum disorders (ASD)^{51,52} (Table 2). The association with suicide risk reached high-quality level based on AMSTAR2, while the two associations with ASD reached moderate quality.

Highly suggestive evidence

For 11 associations (9.2%), there was highly suggestive evidence for the relationship between any antidepressant and increased risk of adverse health outcomes (Table 2), namely attention-deficit hyperactivity disorder (ADHD) in children, cataract development (TCAs), severe bleeding at any site, upper GI bleeding, postpartum hemorrhage, preterm birth, lower APGAR score at 5 minutes, osteoporotic fractures (one for TCAs and one for SSRIs), and risk of hip fracture. Seven of these associations reached moderate quality level based on AMSTAR2 (Table 2). One association with

highly suggestive evidence, however, showed a decreased risk (i.e., protective association) of suicide attempt/completion in adults,⁸⁰ meeting a high-quality level based on AMSTAR2.

Suggestive, weak, and no evidence

There was suggestive evidence for 21 further associations (17.5%) linked to increased risk of adverse health outcomes (eTable 3). For the remaining associations, there was either weak (n=39 [32.5%]) or no evidence (n=46 [38.3%]; all associations with $p>0.05$) (eTables 4 and 5).

Sensitivity analyses

A sensitivity analysis limited to cohort studies, prospective cohort studies, studies controlled for confounding by indication, and North American studies showed that none of the associations within convincing evidence (class I) retained the same rank (Table 3). The most important change was within prospective cohort studies, with one being upgraded to convincing evidence (preterm birth related to any antidepressant).

Another association was upgraded to convincing evidence (lower APGAR scores at 5 minutes) when the sensitivity analysis was limited to SSRIs, as well as the association with preterm birth was also upgraded to convincing evidence when the analysis was limited to other or mixed antidepressants (Table 3).

A further sensitivity analysis limited to non-selected meta-analyses due to overlap agreed with the results of the main analysis (Supplement results and eTable 6). The results of each sensitivity analysis are presented in detail in the Supplement with the full list of covariates in eTable 7.

Discussion

We reviewed 45 meta-analyses of observational studies and we found that only a few of the 74 statistically significant associations between antidepressants and adverse health outcomes had convincing evidence in main and sensitivity analysis, namely the association between antidepressants and increased suicide behavior/completion in individuals aged <19 years old (SSRI),⁸⁰ autism risk in the offspring,^{51,52} preterm birth,⁶⁶ and neonatal adaptation.⁷¹ However, the few convincing

associations do not reflect necessarily causality and, importantly, none of them remained at convincing evidence after accounting for confounding by indication. Overall, our results show that the association between antidepressants and adverse health outcomes is not supported by robust evidence and the underlying disease likely affects the results and inflates significant findings in a relevant way.^{39,44}

To our knowledge, this is the first umbrella review that systematically assesses the potential risk of adverse health outcomes related to antidepressants across a large spectrum of published meta-analyses of observational studies, grading also the evidence by applying well-recognized criteria of its credibility.^{22,23,32,33,36,37} This umbrella review approach has been previously applied to other fields of medicine, such as adverse health outcomes of dietary fiber consumption,³⁷ serum uric acid,²³ and vitamin D.²² Our approach fits to a research field that is undeniably complex and uncertain, as conveyed here.^{22,23,32,33,36,37} The large median number of participants and cases per association allowed to make robust classifications; the number of cases was greater than 1,000 for 87 out of the total of 120 associations. Quality ratings of the included meta-analyses with the AMSTAR2 tool also allowed to interpret our results with confidence. Sensitivity analyses provided additional evidence from cohort, high-quality studies, and studies that were controlled for a psychiatric condition, further increasing reliability of the results.

Our results need to be considered when contemplating the use of antidepressants in children/adolescents integrating this knowledge with data on efficacy from RCTs. A network meta-analysis of RCTs in children/adolescents showed that no antidepressant was superior to placebo apart from fluoxetine, several antidepressants had higher discontinuation rates than placebo, and that venlafaxine increased the risk of suicidality even in the short-term duration of an RCT.¹³ Merging lack of efficacy with lack of safety may raise serious concerns on the use of antidepressants in children/adolescents, with an unfavorable benefit/risk ratio. However, lack of superiority of antidepressants over placebo,¹² especially in children, was due to a very high placebo response,

which has been an increasing problem in RCTs in psychiatry. Also, it is possible that increased suicidality in children/adolescents taking antidepressants may be due to ineffectiveness on depressive symptoms in suicidal individuals, rather than to a direct effect of antidepressants. Additionally, our results show that confounding by indication probably drives safety concerns on the use of antidepressants in children/adolescents. Besides, the risk-benefit evaluation in children and adolescents is quite different for antidepressants (predominantly SSRIs) when used for other disorders like anxiety disorders and OCD.^{3-5,12}

Conversely, we found highly suggestive evidence supporting a protective role of antidepressants in adults against suicidality,⁸⁰ which is consistent with results of a network meta-analysis of RCTs in adults showing that all antidepressants were superior to placebo in reducing depressive symptoms.¹⁰ Similarly, meta-analyses support the efficacy of antidepressants for anxiety disorders⁵ and OCD³ in adults. In adults the benefit/risk ratio must account for clear efficacy of antidepressants and protection against suicide, which should be balanced with other safety concerns rising from the present umbrella review. Overall, several adverse outcomes associated with antidepressants supported by highly suggestive evidence can medically be prevented (i.e., poor bone status, gastro-intestinal bleeding) as previously reported.⁸² Hence, benefits of antidepressants in adults and older adults may well trump (preventable) safety issues, given their clear efficacy in treating various psychiatric disorders.^{3-5,12} Moreover, the association between antidepressants and certain adverse health outcomes varies within specific age groups. For instance, the increased risk of fractures applies predominantly to an older population that is already prone to poor bone status and multimorbidity,⁸³ and not in people aged 20-40 years old.

Convincing evidence before accounting for confounding by indication supporting the association between antidepressants and autism, and other offspring adverse health outcomes may call for restriction of antidepressants during pregnancy to those women with high risk of relapse and with severe clinical presentations. Actually, warnings to avoid prescribing medications in early

pregnancy have been already set.⁸⁴ However, autism remains a rare event, with a prevalence from cohort studies of <1% according to data pooled in this study. Importantly, the convincing level of evidence was not confirmed when considering confounding by indication, suggesting that the association between antidepressants and autism, suicidality in youth, etc. may be due to effects of the underlying disease, rather than to antidepressants,^{39,43,44,50} as recently shown in an umbrella review on risk factors for autism.⁸⁵ Comparing two depression-matched groups with and without antidepressant use may be methodologically more solid than adjusting analyses statistically. In addition, several adverse outcomes had small effect sizes on top of low prevalence and of no proof of a causal relationship between antidepressants and adverse health outcomes.

Hence, given that a depressive episode itself can impair adolescents and both maternal and fetal health, individualized and shared clinical decisions about the benefit-risk ratio of antidepressants during adolescence and pregnancy should be implemented, but adolescence and pregnancy should not be considered absolute contraindications to the use of antidepressants.

Limitations

Several limitations should be mentioned. First, we did not grade the evidence from meta-analyses of RCTs, covering a portion of available studies. However, evidence from RCTs is limited by the selection of healthier patients and frequent short-term follow-up, among other factors.¹⁶ Many severe adverse outcomes cannot be addressed in RCTs, and for low frequency and long-term health risks, observational research is the most feasible method.¹⁷ Nevertheless, observational research is not free from bias, either.^{18,86} Results from observational studies yield associations, which do not imply causality. Further, results in main analyses are affected by various confounders due to lack of randomization, potential channeling bias, and confounding by indication.^{17,18,80} Specifically, the nature of the control groups was only insufficiently characterized, and there is evidence that when matching (and not adjusting) for the underlying psychiatric disorder, risk differences become smaller or non-significant.^{43,44,51} Thus, the association with suicidality may be driven by antidepressants' limited

efficacy in those suicidal children and adolescents according to results from RCTs,¹⁹ rather than being a direct effect of antidepressants on increasing suicidality. Notably, the association between ASD and SSRIs during pregnancy⁵² included studies that were not adjusted for confounders (in contrast with weak evidence of any association between any antidepressants during pregnancy and ASD when adjusting for confounders⁵⁰ – eTable 4). Moreover, no inference can be made about newer antidepressants (e.g., vortioxetine), which have not been assessed in any of the included meta-analyses. Furthermore, there was insufficient data on cardiometabolic outcomes; an emerging concern of the increased prescribing rates of antidepressants, and crucial area for future research.⁸⁷ Finally, our grading system can only provide warnings of the potential presence of systematic biases but cannot provide evidence of their nature and extent^{16,32,33} alongside umbrella reviews cannot supply any comparative ranking, as in network meta-analyses.

Conclusions

Our findings are important in the context of increased antidepressant use globally.^{1,2} An association between antidepressants and few adverse health outcomes with convincing evidence was observed, yet prevalence of those outcomes was on average low (range 0.1%-9.7%), and after accounting for confounding by indication, no association was supported by convincing evidence. Further research in both RCT and real-world samples matched for underlying disease is needed to confirm a possible causal association between antidepressants and adverse outcomes, considering dose-effect response, mechanistic processes, accounting for patient-specific characteristics, such as age, clinical diagnoses, and severity of clinical condition. No absolute contraindication against the use of antidepressants is currently supported by convincing evidence.

Author Contributions

ED, MS had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

ED and MS and EE designed the project.

ED and MS searched the literature, extracted the data, ran the analysis, and contributed equally to this work.

All authors (ED, MS AF, PF-P, PD, TT, BS, JF, MF, DT, AFC, EV, PMcG, AHY, JIS, CUC, EE) approved the protocol, and drafted the manuscript, provided critical comments to the paper for important intellectual content, and approved the final version.

Conflict of Interest Disclosures

Dr. Vieta has received grants and served as consultant, advisor or CME speaker for the following entities: AB-Biotics, Abbott, Allergan, Angelini, AstraZeneca, Bristol-Myers Squibb, Dainippon Sumitomo Pharma, Farindustria, Ferrer, Forest Research Institute, Galenica, Gedeon Richter, Glaxo-Smith-Kline, Janssen, Lundbeck, Otsuka, Pfizer, Roche, SAGE, Sanofi-Aventis, Servier, Shire, Sunovion, Takeda, the Brain and Behaviour Foundation, the Spanish Ministry of Science and Innovation (CIBERSAM), the Seventh European Framework Programme and Horizon 2020, and the Stanley Medical Research Institute.

Dr. Correll has been a consultant and/or advisor to or has received honoraria from: Alkermes, Allergan, Angelini, Boehringer-Ingelheim, Gerson Lehrman Group, Indivior, IntraCellular Therapies, Janssen/J&J, LB Pharma, Lundbeck, MedAvante-ProPhase, Medscape, Merck, Neurocrine, Noven, Otsuka, Pfizer, Rovi, Servier, Sunovion, Supernus, Takeda, and Teva. He has provided expert testimony for Bristol-Myers Squibb, Janssen, and Otsuka. He served on a Data Safety Monitoring Board for Boehringer-Ingelheim, Lundbeck, Rovi, Supernus, and Teva. He received royalties from UpToDate and grant support from Janssen and Takeda. He is also a shareholder of LB Pharma.

Dr. Young is employed by King's College London; Honorary Consultant SLaM (NHS UK). He gave paid lectures and advisory boards for the following companies with drugs used in affective and

related disorders: Astrazenaca, Eli Lilly, Lundbeck, Sunovion, Servier, Livanova, Janssen.

Consultant to Johnson & Johnson. He does not own share holdings in pharmaceutical companies. He has been lead Investigator for Embolden Study (AZ), BCI Neuroplasticity study and Aripiprazole Mania Study. He participated in investigator-initiated studies from AZ, Eli Lilly, Lundbeck, Wyeth, Janssen. He received grant funding (past and present): NIMH (USA); CIHR (Canada); NARSAD (USA); Stanley Medical Research Institute (USA); MRC (UK); Wellcome Trust (UK); Royal College of Physicians (Edin); BMA (UK); UBC-VGH Foundation (Canada); WEDC (Canada); CCS Depression Research Fund (Canada); MSFHR (Canada); NIHR (UK). Janssen (UK)

Paolo Fusar-Poli has been an advisor to Lundbeck outside from this work.

Dr. Firth is supported by a Blackmores Institute Fellowship.

Other authors have no conflicts of interest to declare.

Funding/Support

This study received no funding.

Disclaimer

Brendon Stubbs holds a Clinical Lectureship supported by Health Education England and the NIHR Integrated Clinical Academic (ICA) Programme (ICA-CL-2017-03-001). Brendon Stubbs is also part supported by the Maudsley Charity and the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care South London at King's College Hospital NHS Foundation Trust. The views expressed are those of the author[s] and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Dr. Young states that this report represents independent research funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS

Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health.

Data sharing

The data are available from the corresponding author upon reasonable request.

REFERENCES

1. Pratt LA, Brody DJ, Gu Q. Antidepressant use in persons aged 12 and over: United States, 2005-2008. *NCHS Data Brief*. 2011(76):1-8.
2. Davey CG, Chanen AM. The unfulfilled promise of the antidepressant medications. *Med J Aust*. 2016;204(9):348-350.
3. Skapinakis P, Caldwell DM, Hollingworth W, et al. Pharmacological and psychotherapeutic interventions for management of obsessive-compulsive disorder in adults: a systematic review and network meta-analysis. *Lancet Psychiatry*. 2016;3(8):730-739.
4. Locher C, Koechlin H, Zion SR, et al. Efficacy and Safety of Selective Serotonin Reuptake Inhibitors, Serotonin-Norepinephrine Reuptake Inhibitors, and Placebo for Common Psychiatric Disorders Among Children and Adolescents: A Systematic Review and Meta-analysis. *JAMA Psychiatry*. 2017;74(10):1011-1020.
5. Bighelli I, Castellazzi M, Cipriani A, et al. Antidepressants versus placebo for panic disorder in adults. *Cochrane Database Syst Rev*. 2018;4:CD010676.
6. Leon AC. The revised warning for antidepressants and suicidality: unveiling the black box of statistical analyses. *Am J Psychiatry*. 2007;164(12):1786-1789.
7. Hammad TA, Laughren T, Racoosin J. Suicidality in pediatric patients treated with antidepressant drugs. *Arch Gen Psychiatry*. 2006;63(3):332-339.
8. Gibbons RD, Brown CH, Hur K, Davis J, Mann JJ. Suicidal thoughts and behavior with antidepressant treatment: reanalysis of the randomized placebo-controlled studies of fluoxetine and venlafaxine. *Arch Gen Psychiatry*. 2012;69(6):580-587.

9. Carvalho AF, Sharma MS, Brunoni AR, Vieta E, Fava GA. The Safety, Tolerability and Risks Associated with the Use of Newer Generation Antidepressant Drugs: A Critical Review of the Literature. *Psychother Psychosom.* 2016;85(5):270-288.
10. Cipriani A, Furukawa TA, Salanti G, et al. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. *Lancet.* 2018;391(10128):1357-1366.
11. Gartlehner G, Gaynes BN, Amick HR, et al. Comparative Benefits and Harms of Antidepressant, Psychological, Complementary, and Exercise Treatments for Major Depression: An Evidence Report for a Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med.* 2016;164(5):331-341.
12. Bridge JA, Iyengar S, Salary CB, et al. Clinical response and risk for reported suicidal ideation and suicide attempts in pediatric antidepressant treatment: a meta-analysis of randomized controlled trials. *JAMA.* 2007;297(15):1683-1696.
13. Cipriani A, Zhou X, Del Giovane C, et al. Comparative efficacy and tolerability of antidepressants for major depressive disorder in children and adolescents: a network meta-analysis. *Lancet.* 2016;388(10047):881-890.
14. Moret C, Isaac M, Briley M. Problems associated with long-term treatment with selective serotonin reuptake inhibitors. *J Psychopharmacol.* 2009;23(8):967-974.
15. Corrigan-Curay J, Sacks L, Woodcock J. Real-World Evidence and Real-World Data for Evaluating Drug Safety and Effectiveness. *JAMA.* 2018;320(9):867-868.
16. Solmi M, Correll CU, Carvalho AF, Ioannidis JPA. The role of meta-analyses and umbrella reviews in assessing the harms of psychotropic medications: beyond qualitative synthesis. *Epidemiol Psychiatr Sci.* 2018;27(6):537-542.
17. Benson K, Hartz AJ. A comparison of observational studies and randomized, controlled trials. *N Engl J Med.* 2000;342(25):1878-1886.

18. Kyriacou DN, Lewis RJ. Confounding by Indication in Clinical Research. *JAMA*. 2016;316(17):1818-1819.
19. Cipriani A, Barbui C, Geddes JR. Suicide, depression, and antidepressants. *BMJ*. 2005;330(7488):373-374.
20. Ioannidis JP. Integration of evidence from multiple meta-analyses: a primer on umbrella reviews, treatment networks and multiple treatments meta-analyses. *CMAJ*. 2009;181(8):488-493.
21. Ioannidis J. Next-generation systematic reviews: prospective meta-analysis, individual-level data, networks and umbrella reviews. *Br J Sports Med*. 2017;51(20):1456-1458.
22. Theodoratou E, Tzoulaki I, Zgaga L, Ioannidis JP. Vitamin D and multiple health outcomes: umbrella review of systematic reviews and meta-analyses of observational studies and randomised trials. *BMJ*. 2014;348:g2035.
23. Li X, Meng X, Timofeeva M, et al. Serum uric acid levels and multiple health outcomes: umbrella review of evidence from observational studies, randomised controlled trials, and Mendelian randomisation studies. *BMJ*. 2017;357:j2376.
24. Radua J, Ramella-Cravaro V, Ioannidis JPA, et al. What causes psychosis? An umbrella review of risk and protective factors. *World Psychiatry*. 2018;17(1):49-66.
25. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
26. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA*. 2000;283(15):2008-2012.
27. Raglan O, Kalliala I, Markozannes G, et al. Risk factors for endometrial cancer: An umbrella review of the literature. *Int J Cancer*. 2018.
28. Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017;358:j4008.

29. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3):177-188.
30. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327(7414):557-560.
31. Riley RD, Higgins JP, Deeks JJ. Interpretation of random effects meta-analyses. *BMJ*. 2011;342:d549.
32. Bortolato B, Kohler CA, Evangelou E, et al. Systematic assessment of environmental risk factors for bipolar disorder: an umbrella review of systematic reviews and meta-analyses. *Bipolar Disord*. 2017;19(2):84-96.
33. Dragioti E, Evangelou E, Larsson B, Gerdle B. Effectiveness of multidisciplinary programmes for clinical pain conditions: An umbrella review. *J Rehabil Med*. 2018.
34. Ioannidis JP, Trikalinos TA. An exploratory test for an excess of significant findings. *Clin Trials*. 2007;4(3):245-253.
35. Ioannidis JPA. Clarifications on the application and interpretation of the test for excess significance and its extensions. *Journal of Mathematical Psychology*. 2013;57(5):184-187.
36. Dragioti E, Karathanos V, Gerdle B, Evangelou E. Does psychotherapy work? An umbrella review of meta-analyses of randomized controlled trials. *Acta Psychiatr Scand*. 2017;136(3):236-246.
37. Veronese N, Solmi M, Caruso MG, et al. Dietary fiber and health outcomes: an umbrella review of systematic reviews and meta-analyses. *Am J Clin Nutr*. 2018;107(3):436-444.

Barbui C, Esposito E, Cipriani A. Selective serotonin reuptake inhibitors and risk of suicide: a systematic review of observational studies. *CMAJ*. 2009;180(3):291-297.

Morales DR, Slattery J, Evans S, Kurz X. Antidepressant use during pregnancy and risk of autism spectrum disorder and attention deficit hyperactivity disorder: systematic review of observational studies and methodological considerations. *BMC Med*. 2018;16(1):6.

Andalib S, Emamhadi MR, Yousefzadeh-Chabok S, et al. Maternal SSRI exposure increases the risk of autistic offspring: A meta-analysis and systematic review. *Eur Psychiatry*. 2017;45:161-166.

- Huang H, Coleman S, Bridge JA, Yonkers K, Katon W. A meta-analysis of the relationship between antidepressant use in pregnancy and the risk of preterm birth and low birth weight. *Gen Hosp Psychiatry*. 2014;36(1):13-18.
- Ross LE, Grigoriadis S, Mamisashvili L, et al. Selected pregnancy and delivery outcomes after exposure to antidepressant medication: a systematic review and meta-analysis. *JAMA Psychiatry*. 2013;70(4):436-443.
38. Masarwa R, Bar-Oz B, Gorelik E, Reif S, Perlman A, Matok I. Prenatal exposure to selective serotonin reuptake inhibitors and serotonin norepinephrine reuptake inhibitors and risk for persistent pulmonary hypertension of the newborn: a systematic review, meta-analysis, and network meta-analysis. *Am J Obstet Gynecol*. 2019 ;220(1): 57.e1-57.e13.
39. Halvorsen A, Hesel B, Ostergaard SD, Danielsen AA. In Utero Exposure to SSRIs and Development of Mental Disorders: A Systematic Review and Meta-analysis. *Acta Psychiatr Scand*. 2019 ;139(6):493-507.
40. Wang YC, Tai PA, Poly TN, et al. Increased Risk of Dementia in Patients with Antidepressants: A Meta-Analysis of Observational Studies. *Behav Neurol*. 2018 ;2018:5315098.
41. Schweiger JU, Schweiger U, Hüppe M, et al. The Use of Antidepressive Agents and Bone Mineral Density in Women: A Meta-Analysis. *Int J Environ Res Public Health*. 2018;15(7): pii: E1373.
42. Khanassov V, Hu J, Reeves D, van Marwijk H. Selective serotonin reuptake inhibitor and selective serotonin and norepinephrine reuptake inhibitor use and risk of fractures in adults: A systematic review and meta-analysis. *Int J Geriatr Psychiatry*. 2018 ;33(12):1688-1708.
43. Jiang HY, Peng CT, Zhang X, Ruan B. Antidepressant use during pregnancy and the risk of attention-deficit/hyperactivity disorder in the children: a meta-analysis of cohort studies. *BJOG*. 2018;125(9):1077-1084.
44. Gao SY, Wu QJ, Sun C, et al. Selective serotonin reuptake inhibitor use during early pregnancy and congenital malformations: a systematic review and meta-analysis of cohort studies of more than 9 million births. *BMC Med*. 2018;16(1):205.

45. Guan HB, Wei Y, Wang LL, Qiao C, Liu CX. Prenatal Selective Serotonin Reuptake Inhibitor Use and Associated Risk for Gestational Hypertension and Preeclampsia: A Meta-Analysis of Cohort Studies. *J Womens Health (Larchmt)*. 2018;27(6):791-800.
46. Chappuis V, Avila-Ortiz G, Araújo MG, Monje A. Medication-related dental implant failure: Systematic review and meta-analysis. *Clin Oral Implants Res*. 2018;29 (Suppl 16):55-68.
47. Na KS, Jung HY, Cho SJ, Cho SE. Can we recommend mirtazapine and bupropion for patients at risk for bleeding?: A systematic review and meta-analysis. *J Affect Disord*. 2018;225:221-226.
48. Man KKC, Chan EW, Ip P, et al. Prenatal antidepressant exposure and the risk of attention-deficit hyperactivity disorder in children: A systematic review and meta-analysis. *Neurosci Biobehav Rev*. 2018;86:1-11.
49. Fu YD, Q.; Zhu, L.; Wu, S. Antidepressants use and risk of cataract development: a systematic review and meta-analysis. *BMC ophthalmology*. 2018;18(1):31.
50. Zhou XH, Li YJ, Ou JJ, Li YM. Association between maternal antidepressant use during pregnancy and autism spectrum disorder: an updated meta-analysis. *Mol Autism*. 2018;9:21.
51. Morales DR, Slattery J, Evans S, Kurz X. Antidepressant use during pregnancy and risk of autism spectrum disorder and attention deficit hyperactivity disorder: systematic review of observational studies and methodological considerations. *BMC Med*. 2018;16(1):6.
52. Andalib S, Emamhadi MR, Yousefzadeh-Chabok S, et al. Maternal SSRI exposure increases the risk of autistic offspring: A meta-analysis and systematic review. *Eur Psychiatry*. 2017;45:161-166.
53. Zhang TN, Gao SY, Shen ZQ, et al. Use of selective serotonin-reuptake inhibitors in the first trimester and risk of cardiovascular-related malformations: a meta-analysis of cohort studies. *Sci Rep*. 2017;7:43085.
54. Shen ZQ, Gao SY, Li SX, et al. Sertraline use in the first trimester and risk of congenital anomalies: a systemic review and meta-analysis of cohort studies. *Br J Clin Pharmacol*. 2017;83(4):909-922.
55. Laporte S, Chapelle C, Caillet P, et al. Bleeding risk under selective serotonin reuptake inhibitor (SSRI) antidepressants: A meta-analysis of observational studies. *Pharmacol Res*. 2017;118:19-32.

56. Gao SY, Wu QJ, Zhang TN, et al. Fluoxetine and congenital malformations: a systematic review and meta-analysis of cohort studies. *Br J Clin Pharmacol*. 2017;83(10):2134-2147.
57. Biffi A, Scotti L, Corrao G. Use of antidepressants and the risk of cardiovascular and cerebrovascular disease: a meta-analysis of observational studies. *Eur J Clin Pharmacol*. 2017;73(4):487-497.
58. Jiang HY, Xu LL, Li YC, Deng M, Peng CT, Ruan B. Antidepressant use during pregnancy and risk of postpartum hemorrhage: A systematic review and meta-analysis. *J Psychiatr Res*. 2016;83:160-167.
59. Healy D, Le Noury J, Mangin D. Links between serotonin reuptake inhibition during pregnancy and neurodevelopmental delay/spectrum disorders: A systematic review of epidemiological and physiological evidence. *Int J Risk Saf Med*. 2016;28(3):125-141.
60. Eke AC, Saccone G, Berghella V. Selective serotonin reuptake inhibitor (SSRI) use during pregnancy and risk of preterm birth: a systematic review and meta-analysis. *BJOG*. 2016;123(12):1900-1907.
61. Bérard A, Iessa N, Chaabane S, Muanda FT, Boukhris T, Zhao JP. The risk of major cardiac malformations associated with paroxetine use during the first trimester of pregnancy: a systematic review and meta-analysis. *Br J Clin Pharmacol*. 2016;81(4):589-604.
62. Undela K, Parthasarathi G, John SS. Impact of antidepressants use on risk of myocardial infarction: A systematic review and meta-analysis. *Indian J Pharmacol*. 2015;47(3):256-262.
63. Singh I, Achuthan S, Chakrabarti A, Rajagopalan S, Srinivasan A, Hota D. Influence of pre-operative use of serotonergic antidepressants (SADs) on the risk of bleeding in patients undergoing different surgical interventions: a meta-analysis. *Pharmacoepidemiol Drug Saf*. 2015;24(3):237-245.
64. Jiang HY, Chen HZ, Hu XJ, et al. Use of selective serotonin reuptake inhibitors and risk of upper gastrointestinal bleeding: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol*. 2015;13(1):42-50.
65. Shin D, Oh YH, Eom CS, Park SM. Use of selective serotonin reuptake inhibitors and risk of stroke: a systematic review and meta-analysis. *J Neurol*. 2014;261(4):686-695.

66. Huang H, Coleman S, Bridge JA, Yonkers K, Katon W. A meta-analysis of the relationship between antidepressant use in pregnancy and the risk of preterm birth and low birth weight. *Gen Hosp Psychiatry*. 2014 ;36(1):13-18.
67. Huybrechts KF, Sanghani RS, Avorn J, Urato AC. Preterm birth and antidepressant medication use during pregnancy: a systematic review and meta-analysis. *PloS One*. 2014;9(3):e92778.
68. Grigoriadis S, Vonderporten EH, Mamisashvili L, et al. Prenatal exposure to antidepressants and persistent pulmonary hypertension of the newborn: systematic review and meta-analysis. *BMJ*. 2014;348:f6932.
69. Oh SW, Kim J, Myung SK, Hwang SS, Yoon DH. Antidepressant use and risk of coronary heart disease: meta-analysis of observational studies. *Br J Clin Pharmacol*. 2014;78(4):727-737.
70. Wu Q, Qu W, Crowell MD, Hentz JG, Frey KA. Tricyclic antidepressant use and risk of fractures: a meta-analysis of cohort and case-control studies. *J Bone Miner Res*. 2013;28(4):753-763.
71. Ross LE, Grigoriadis S, Mamisashvili L, et al. Selected pregnancy and delivery outcomes after exposure to antidepressant medication: a systematic review and meta-analysis. *JAMA Psychiatry*. 2013;70(4):436-443.
72. Riggall L, Frankel Z, Moretti M, Pupco A, Koren G. The fetal safety of fluoxetine: a systematic review and meta-analysis. *J Obstet Gynaecol Can*. 2013;35(4):362-369.
73. Grigoriadis S, VonderPorten EH, Mamisashvili L, et al. The effect of prenatal antidepressant exposure on neonatal adaptation: a systematic review and meta-analysis. *J Clin Psychiatry*. 2013;74(4):e309-320.
74. Myles N, Newall H, Ward H, Large M. Systematic meta-analysis of individual selective serotonin reuptake inhibitor medications and congenital malformations. *Aust N Z J Psychiatry*. 2013;47(11):1002-1012.
75. Nikfar S, Rahimi R, Hendoiee N, Abdollahi M. Increasing the risk of spontaneous abortion and major malformations in newborns following use of serotonin reuptake inhibitors during pregnancy: A systematic review and updated meta-analysis. *Daru*. 2012;20(1):75.

76. Oderda LH, Young JR, Asche CV, Pepper GA. Psychotropic-related hip fractures: meta-analysis of first-generation and second-generation antidepressant and antipsychotic drugs. *Ann Pharmacother*. 2012;46(7-8):917-28.
77. Eom CS, Park SM, Cho KH. Use of antidepressants and the risk of breast cancer: a meta-analysis. *Breast Cancer Res Treat*. 2012;136(3):635-645.
78. Lee HK, Eom CS, Kwon YM, Ahn JS, Kim S, Park SM. Meta-analysis: selective serotonin reuptake inhibitors and colon cancer. *Eur J Gastroenterol Hepatol*. 2012;24(10):1153-1157.
79. Cosgrove L, Shi L, Creasey DE, Anaya-McKivergan M, Myers JA, Huybrechts KF. Antidepressants and breast and ovarian cancer risk: a review of the literature and researchers' financial associations with industry. *PLoS One*. 2011;6(4):e18210.
80. Barbui C, Esposito E, Cipriani A. Selective serotonin reuptake inhibitors and risk of suicide: a systematic review of observational studies. *CMAJ*. 2009;180(3):291-297.
81. Hemels ME, Einarson A, Koren G, Lanctot KL, Einarson TR. Antidepressant use during pregnancy and the rates of spontaneous abortions: a meta-analysis. *Ann Pharmacother*. 2005;39(5):803-809.
82. National Institute for Health and Care Excellence. *Depression in adults*. Clinical guideline (CG 90). <http://www.nice.org.uk/> Published Oct 2009. Accessed November 2018.
83. Brannstrom J, Lovheim H, Gustafson Y, Nordstrom P. Association Between Antidepressant Drug Use and Hip Fracture in Older People Before and After Treatment Initiation. *JAMA Psychiatry*. 2019;76(2):172-179.
84. British Medical Association; Royal Pharmaceutical Society. *British National Formulary (BNF)* 57. London, UK: BMJ Group and RPS Publishing; March 2009.
85. Yeob Kim J, Ji Son M, Yun Son C, et al. Environmental risk factors and biomarkers for autism spectrum disorder: an umbrella review of the evidence. *The Lancet Psychiatry*. 2019;in press.
86. Howland RH. Antidepressant medication and pregnancy: time for randomized controlled trials. *J Psychosoc Nurs Ment Health Serv*. 2013;51(2):11-14.

- 87.. Firth J, Siddiqi N, Koyanagi A, et al. A Blueprint for Protecting Physical Health in People with Mental Illness: Directions for Health Promotion, Clinical Services and Future Research. *The Lancet Psychiatry*. 2019;in press.

Tables

Table 1 Criteria for evaluation of the credibility of the evidence of observational studies

Classification	Criteria
Convincing evidence (Class I)	<ol style="list-style-type: none"> 1. More than 1000 cases 2. Significant summary associations ($p < 1 \times 10^{-6}$) per random-effects calculations 3. No evidence of small-study effects 4. No evidence of excess of significance bias 5. Prediction intervals not including the null value 6. Largest study nominally significant ($p < 0.05$) 7. No large heterogeneity (i.e., $I^2 < 50\%$)
Highly Suggestive evidence (Class II)	<ol style="list-style-type: none"> 1. More than 1000 cases 2. Significant summary associations ($p < 1 \times 10^{-6}$) per random-effects calculation 3. Largest study nominally significant ($p < 0.05$)
Suggestive Evidence (Class III)	<ol style="list-style-type: none"> 1. More than 1000 cases 2. Significant summary associations ($p < 1 \times 10^{-3}$) per random-effects calculations
Weak evidence (Class IV)	<ol style="list-style-type: none"> 1. All other associations with $p \leq 0.05$
Non-significant associations (NS)	<ol style="list-style-type: none"> 1. All associations with $p > 0.05$

Table 2 Convincing evidence (Class I) and highly suggestive evidence (Class II) for the association of antidepressants and risk of adverse health outcomes in meta-analyses of observational studies

Adverse health outcomes (author, year)	Exposed/ Unexposed	Prevalence (%) based on cohort studies	n	Random-effects measure, ES (95% CI)	Results	Criteria used for classification of level of evidence							AMSTAR 2 quality
						N cases/ total population	p- value random effects	I ² (p- value)	PI (95 % CI)	SSE/ES B	LS	CE	
Autism spectrum disorders (Pre-pregnancy maternal exposure; Morales, 2018) ⁵¹	Any AD users/ No AD users	0. 8%	7	RR, 1.48 (1.29, 1.71)	Increased risk for ADs	22 877/ 2 400 720	6.8x10 ⁻⁸	24 (0.244)	1.09–2.02	No/NP	Yes	I	Moderate
Autism spectrum disorders (pregnancy maternal exposure unadjusted estimates only; Andalib, 2017) ⁵²	SSRIs / Non-SSRIs users	0. 9%	7	OR, 1.84 (1.60, 2.11)	Increased risk for SSRIs	58 178 / 5 868 692	1.2x10 ⁻¹⁷	0 (0.729)	1.53–2.20	No/NP	Yes	I	Moderate

						Criteria used for classification of level of evidence							AMSTAR 2 quality
Adverse health outcomes (author, year)	Exposed/ Unexposed	Prevalence (%) based on cohort studies	n	Random- effects measure, ES (95% CI)	Results	N cases/ total population	p- value random effects	I ² (p- value)	PI (95 % CI)	SSE/ES B	LS	CE	
Suicide attempt and completion in children and adolescents (Barbui, 2009) ⁸⁰	SSRIs / Non-SSRIs users	9. 4%	5	OR, 1.92 (1.51, 2.44)	Increased risk for SSRIs	6 531/ 61 522	1.0x10 ⁻⁷	0 (0.469)	1.30–2.84	No/NP	Yes	I	High
Osteoporotic fractures (Khanassov, 2018) ⁴²	SSRIs / Non-SSRIs users	6. 5%	24	RR, 1.67 (1.56, 1.80)	Increased risk for SSRIs	136 449/ 1 546 913	1.3x10 ⁻⁴⁴	88 (<0.000)	1.23-2.27	No/NP	Yes	II	Moderate
Attention-deficit hyperactivity disorder in children (Man, 2018) ⁴⁸	Prenatal exposure to ADs/ No AD users	2. 4%	7	RR, 1.39 (1.21,1.61)	Increased risk for ADs	57 552/ 2 886 904	5.1x10 ⁻⁶	79 (<0.000)	0.90–2.15	No/NP	Yes	II	Moderate
Cataract development (Fu, 2018) ⁴⁹	TCAs/ non-users or no users of any other AD	NA	3	OR, 1.19 (1.11, 1.28)	Increased risk for TCAs	215 298/ 431 171	2.0x10 ⁻⁶	58 (0.092)	0.56-2.52	No/NP	Yes	II	Moderate
Severe bleeding at any site	SSRIs+ SNRIs/ non-users or no	2. 4%	44	OR, 1.41 (1.27,1.57)	Increased risk for	75 215/ 1 443 029	2.2x10 ⁻¹⁰	90 (<0.000)	0.77-2.59	No/No	Yes	II	Low

						Criteria used for classification of level of evidence							AMSTAR 2 quality
Adverse health outcomes (author, year)	Exposed/ Unexposed	Prevalence (%) based on cohort studies	n	Random- effects measure, ES (95% CI)	Results	N cases/ total population	p- value random effects	I ² (p- value)	PI (95 % CI)	SSE/ES B	LS	CE	
(Laporte, 2017) ⁵⁵	users of any other AD				SSRIs+ SNRIs								
Postpartum hemorrhage (Jiang, 2016) ⁵⁸	Any AD users/ No AD users	6. 8%	17	RR, 1.32 (1.17, 1.48)	Increased risk for ADs	49 155/ 65 1715	3.3x10 ⁻⁶	85 (<0.000)	0.84-2.07	No/NP	Yes	II	Low
Upper GI bleeding (Jiang, 2015) ⁶⁴	SSRIs+others non- ADs/ No SSRIs use only+others non- ADs	0.7%	22	OR, 1.55 (1.35, 1.78)	Increased risk for SSRIs	56 182/ 592 508	9.2x10 ⁻¹²	89 (<0.000)	0.83-2.91	No/No	Yes	II	Moderate
Preterm birth (Huang, 2014) ⁶⁶	Any AD users/ No AD users	0. 8%	28	RR, 1.68 (1.52, 1.86)	Increased risk for ADs	24 669/ 3 063709	3.6x10 ⁻²³	44 (0.008)	1.23-2.30	Yes/NP	Yes	II	Moderate
Osteoporotic fractures (Wu, 2013) ⁷⁰	TCAs users/ Non-TCAs users	1. 4%	12	RR, 1.45 (1.31, 1.60)	Increased risk for TCAs	178 237/ 831 912	6.2x10 ⁻¹³	76 (<0.000)	1.04-2.01	Yes/No	Yes	II	Moderate
Apgar score at 5 minutes	Any AD users/ No AD users	2. 1%	15	SMD, -0.33	Increased risk for ADs	1 473/ 71 828	7.5x10 ⁻⁷	58 (0.003)	-1.02, 0.36	No/No	Yes	II	Moderate

Adverse health outcomes (author, year)	Exposed/ Unexposed	Prevalence (%) based on cohort studies	n	Random-effects measure, ES (95% CI)	Results	Criteria used for classification of level of evidence							AMSTAR 2 quality
						N cases/ total population	p- value random effects	I ² (p- value)	PI (95 % CI)	SSE/ES B	LS	CE	
(Ross, 2013) ⁷¹				(-0.47,- 0.20)									
Hip Fracture (Oderda, 2012) ⁷⁶	TCAs and/or SSRIs users/ No AD users	7. 4%	18	OR, 1.78 (1.53, 2.07)	Increased risk for TCAs or SSRIs	49 276/ 210 577	5.2x10 ⁻¹⁴	89 (<0.000)	1.00-3.19	Yes/Yes	Yes	II	Critically low
Suicide attempt and completion in adults (Barbui, 2009) ⁸⁰	SSRIs / Non- SSRIs users	4.5%	7	OR, 0.59 (0.48, 0.72)	Decreased risk for SSRIs	7 164/ 147 383	5.2x10 ⁻⁷	59 (0.023)	0.33-1.05	No/NP	Yes	II	High

AD–antidepressants, APGAR – acronym of Appearance (skin color), Pulse (heart rate), Grimace (reflex irritability), Activity (muscle tone), and Respiration, CE – class of evidence, CI – confidence interval, ES – effect size, ESB –excess significance bias, GI–gastrointestinal, I²–heterogeneity, LS – largest study with significant effect, n – number of included studies per association, N – number of cases, NA– not applicable, NP – not pertinent because the number of observed studies is less than the expected, OR – odds ratio, PI – prediction interval, RR – relative risk, SNRIs – serotonin–norepinephrine reuptake inhibitors, SSRIs – selective serotonin reuptake inhibitors, SMD–standardised mean difference, SSE – small-study effect, TCAs – tricyclic antidepressants.

Table 3 Sensitivity analysis for the associations between antidepressants use and risk of adverse health outcomes of class I-II associations limited to only cohort studies, prospective cohort, studies with adjustments for multiple covariates, studies adjusted for confounding by indication, high-quality studies, classes of antidepressants (SSRIs studies, TCAs studies, other or mixed antidepressants studies), and study location (Europe, North America, other regions)

					Criteria used for classification of level of evidence							
Adverse health outcomes (author, year)	Exposed/Unexposed	Prevalence (%) based on cohort studies	n	Random- effects measure, ES (95% CI)	N cases/ total population	p- random effects	I ² (p)	PI (95% CI)	SSE/ESB	LS	CE	CES / AMSTAR 2
Cohort studies (both retrospective and prospective)												
Autism spectrum disorders (pregnancy maternal exposure; unadjusted estimates only; Andalib, 2017) ⁵²	SSRIs/Non-SSRIs users	0. 9%	3	OR, 1.65 (1.37, 2.00)	50 494 / 5 790 186	2.2x10 ⁻⁷	0 (0.692)	0.48-5.68	No/No	Yes	I	II/ Moderate
Osteoporotic fractures (Khanassov, 2018) ⁴²	SSRIs / Non-SSRIs users	6. 5%	16	RR, 1.63 (1.49, 1.79)	56 397/ 859 611	8.9x10 ⁻²⁷	85 (<0.000)	1.20-2.22	No/NP	Yes	II	II/ Moderate

Preterm birth (Huang, 2014) ⁶⁶	Any AD users/ Non-AD users	0. 8%	24	RR, 1.67 (1.51, 1.86)	24 378/ 3 063 012	6.5x10 ⁻²²	49 (0.004)	1.21-2.33	Yes/No	Yes	II	II / Moderate
Apgar score at 5 minutes (Ross, 2013) ⁷¹	Any AD users/ Non-AD users	2. 1%	15	SMD, -0.33 (-0.47,-0.20)	1 473/ 71 828	7.5x10 ⁻⁷	58 (0.003)	-1.02, 0.36	No/No	Yes	II	II / Moderate
Suicide attempt and completion in children and adolescents (Barbui, 2009) ⁸⁰	SSRIs / Non-SSRIs users	9. 4%	3	OR, 1.89 (1.46, 2.43)	5 647/ 59 971	1.8x10 ⁻⁷	0 (0.463)	0.36-9.85	No/NP	Yes	I	II / High
Suicide attempt and completion in adults (Barbui, 2009) ⁸⁰	SSRIs / Non-SSRIs users	4.5%	5	OR, 0.53 (0.43, 0.66)	6 458/ 143 340	5.2x10 ⁻⁷	55 (0.064)	0.27-1.04	No/NP	Yes	II	II / High
Prospective cohort studies												
Autism spectrum disorders (pregnancy maternal exposure; unadjusted estimates only; Andalib, 2017) ⁵²	SSRIs/Non-SSRIs users	0. 9%	3	OR, 1.65 (1.37, 2.00)	50 494 / 5 790 186	2.2x10 ⁻⁷	0 (0.692)	0.48-5.68	No/No	Yes	I	II/ Moderate
Preterm birth (Huang, 2014) ⁶⁶	Any AD users/ Non-AD users	0. 4%	11	RR, 1.87 (1.52, 2.30)	2 540/ 690 121	3.4x10 ⁻⁹	31 (0.148)	1.17-3.00	No/NP	Yes	II	I / Moderate
Studies with adjustments for multiple covariates												
Autism spectrum disorders (Pre-pregnancy maternal exposure; Morales, 2018) ⁵¹	Any AD users/ No AD users	0. 8%	7	RR, 1.48 (1.29, 1.71)	22 877/ 2 400 720	6.8x10 ⁻⁸	24 (0.244)	1.09-2.02	No/NP	Yes	I	I/ Moderate

Suicide attempt and completion in children and adolescents (Barbui, 2009) ⁸⁰	SSRIs / Non-SSRIs users	9. 4%	5	OR, 1.92 (1.51, 2.44)	6 531/ 61 522	1.0x10 ⁻⁷	0 (0.469)	1.30–2.84	No/NP	Yes	I	I / High
Osteoporotic fractures (Khanassov, 2018) ⁴²	SSRIs / Non-SSRIs users	6.5%	22	RR, 1.67 (1.53, 1.82)	96 353/ 929 936	6.6x10 ⁻³³	88 (<0.000)	1.17-2.38	No/NP	Yes	II	II/ Moderate
Attention-deficit hyperactivity disorder in children (Man, 2018) ⁴⁸	Prenatal exposure to ADs/ No AD users	2. 4%	7	RR, 1.39 (1.21,1.61)	57 552/ 2 886 904	5.1x10 ⁻⁶	79 (<0.000)	0.90–2.15	No/NP	Yes	II	II/ Moderate
Upper GI bleeding (Jiang, 2015) ⁶⁴	SSRIs+others non- ADs/ No SSRIs use only+others non- ADs	1.3%	15	RR, 1.48 (1.32, 1.67)	43 571/ 308 060	1.3x10 ⁻¹⁰	61 (0.001)	1.00-2.20	No/NP	Yes	II	II / Moderate
Osteoporotic fractures (Wu, 2013) ⁷⁰	TCAs users/ Non-TCAs users	NA	11	RR, 1.43 (1.29, 1.58)	178 237 / 740 768	1.9 x10 ⁻¹²	77 (<0.000)	1.04-1.97	Yes/No	Yes	II	II / Moderate
Hip Fracture (Oderda, 2012)	TCAs and/or SSRIs users/ No AD users	NA	14	OR, 1.76 (1.49, 2.08)	47 762/ 198 820	1.9x10 ⁻¹¹	92 (0.001)	0.96-3.24	Yes/No	Yes	II	II / Critically low
Studies adjusted for confounding by indication												
Autism spectrum disorders (Pre-pregnancy maternal exposure; Morales, 2018) ⁵¹	Any AD users/ No AD users	1.1%	3	RR, 1.69 (1.39, 2.05)	3 016/ 667 431	1.0 x10 ⁻⁷	0 (0.735)	0.48–5.94	No/No	Yes	I	II/ Moderate

Suicide attempt and completion in children and adolescents (Barbui, 2009) ⁸⁰	SSRIs / Non-SSRIs users	9.7%	4	OR, 2.04 (1.23, 3.41)	5 522 / 55 124	0.006	16 (0.314)	0.47-8.81	Yes/NP	No	I	IV / High
Osteoporotic fractures (Khanassov, 2018) ⁴²	SSRIs / Non-SSRIs users	6.3%	11	RR, 1.61 (1.43, 1.82)	54 023/ 645 463	1.0x10 ⁻¹⁴	88 (<0.000)	1.08-2.41	No/NP	Yes	II	II/ Moderate
Osteoporotic fractures (Wu, 2013) ⁷⁰	TCAs users/ Non-TCAs users	NA	7	RR, 1.47 (1.23, 1.69)	156 374/ 659 389	5.0x10 ⁻⁸	62 (0.021)	0.98-2.22	No/No	Yes	II	II / Moderate
High-quality primary studies												
Autism spectrum disorders (Pre-pregnancy maternal exposure; Morales, 2018) ⁵¹	Any AD users/ No AD users	0. 8%	7	RR, 1.48 (1.29, 1.71)	22 877/ 2 400 720	6.8x10 ⁻⁸	24 (0.244)	1.09–2.02	No/NP	Yes	I	I/ Moderate
Autism spectrum disorders (pregnancy maternal exposure; unadjusted estimates only; Andalib, 2017) ⁵²	SSRIs / Non-SSRIs users	0. 9%	7	OR, 1.84 (1.60, 2.11)	58 178 / 5 868 692	1.2x10 ⁻¹⁷	0 (0.729)	1.53–2.20	No/NP	Yes	I	I/ Moderate
Suicide attempt and completion in children and adolescents (Barbui, 2009) ⁸⁰	SSRIs / Non-SSRIs users	3.6%	4	OR, 1.88 (1.47, 2.40)	5 961/ 30 343	3.5x10 ⁻⁷	0 (0.496)	1.10-3.21	No/NP	Yes	I	I/ High
Osteoporotic fractures (Khanassov, 2018) ⁴²	SSRIs / Non-SSRIs users	6. 6%	20	RR, 1.70 (1.57, 1.85)	117 567/ 1 053 697	9.6x10 ⁻³⁷	88 (<0.000)	1.23-2.35	No/NP	Yes	II	II/ Moderate
Attention-deficit hyperactivity disorder in children	Prenatal exposure to ADs/ No AD users	2. 4%	7	RR, 1.39 (1.21,1.61)	57 552/ 2 886 904	5.1x10 ⁻⁶	79 (<0.000)	0.90–2.15	No/NP	Yes	II	II/ Moderate

(Man, 2018) ⁴⁸												
Postpartum hemorrhage (Jiang, 2016) ⁵⁸	Any AD users/ No AD users	6. 8%	16	RR, 1.32 (1.17, 1.48)	49 142/ 651 439	3.7x10 ⁻⁶	86 (<0.000)	0.84-2.08	No/NP	Yes	II	II Low
Preterm birth (Huang, 2014) ⁶⁶	Any AD users/ No AD users	0. 3%	19	RR, 1.86 (1.59, 2.19)	5 116/ 1 658 666	5.3x10 ⁻¹⁴	52 (0.004)	1.16-3.00	Yes/No	Yes	II	II/ Moderate
Osteoporotic fractures (Wu, 2013) ⁷⁰	TCAs users/ Non-TCAs users	NA	9	RR, 1.46 (1.30, 1.65)	36 865/ 247 078	6.0x10 ⁻¹⁰	82 (<0.000)	0.99-2.15	Yes /NP	Yes	II	II/ Moderate
Hip Fracture (Oderda, 2012) ⁷⁶	TCAs and/or SSRIs users/ No AD users	NA	6	OR, 1.59 (1.31, 1.92)	41 227/ 159 831	2.5x10 ⁻⁷	91 (<0.000)	0.82-3.07	Yes/No	Yes	II	II/ Critically low
Suicide attempt and completion in adults (Barbui, 2009) ⁸⁰	SSRIs / Non- SSRIs users	4.5%	7	OR, 0.59 (0.48, 0.72)	7 164/ 147 383	5.2x10 ⁻⁷	59 (0.023)	0.33-1.05	No/NP	Yes	II	II/ High
SSRIs studies												
Autism spectrum disorders (pregnancy maternal exposure; unadjusted estimates only; Andalib, 2017) ⁵²	SSRIs / Non-SSRIs users	0. 9%	7	OR, 1.84 (1.60, 2.11)	58 178 / 5 868 692	1.2x10 ⁻¹⁷	0 (0.729)	1.53–2.20	No/NP	Yes	I	I/ Moderate
Suicide attempt and completion in children and adolescents (Barbui, 2009) ⁸⁰	SSRIs / Non-SSRIs users	9. 4%	5	OR, 1.92 (1.51, 2.44)	6 531/ 61 522	1.0x10 ⁻⁷	0 (0.469)	1.30–2.84	No/NP	Yes	I	I / High
Apgar score at 5 minutes (Ross, 2013) ⁷¹	SSRIs / Non-SSRIs users	5. 7%	13	SMD, -0.27 (-0.37, -0.16)	1 127/ 19 695	2.1x10 ⁻⁷	35 (0.100)	-0.53, -0.01	No/No	Yes	II	I / Moderate

Osteoporotic fractures (Khanassov, 2018) ⁴²	SSRIs / Non-SSRIs users	6. 5%	24	RR, 1.67 (1.56, 1.80)	136 449/ 1 546 913	1.3x10 ⁻⁴⁴	88 (<0.000)	1.23-2.27	No/NP	Yes	II	II/ Moderate
Preterm birth (Huang, 2014) ⁶⁶	SSRIs / Non-SSRIs users	9. 7%	20	RR, 1.73 (1.53, 1.96)	21 163/ 2 153 680	3.6x10 ⁻²³	48 (0.009)	1.22-2.46	Yes/NP	Yes	II	II / Moderate
Suicide attempt and completion in adults (Barbui, 2009) ⁸⁰	SSRIs / Non- SSRIs users	3. 5%	7	OR, 0.59 (0.48, 0.72)	7 164/ 147 383	5.2x10 ⁻⁷	59 (0.023)	0.33-1.05	No/NP	Yes	II	II / High
TCAs studies												
Cataract development (Fu, 2018) ⁴⁹	TCAs/ non-users or no users of any other AD	NA	3	OR, 1.19 (1.11, 1.28)	215 298/ 431 171	2.0x10 ⁻⁶	58 (0.092)	0.56-2.52	No/NP	Yes	II	II / Moderate
Osteoporotic fractures (Wu, 2013) ⁷⁰	TCA users/ Non- TCA users	1. 4%	12	RR, 1.45 (1.31, 1.60)	178 237/ 831 912	6.2x10 ⁻¹³	76 (<0.000)	1.04-2.01	Yes/No	Yes	II	II / Moderate
Other or mixed AD studies												
Autism spectrum disorders (Pre-pregnancy maternal exposure; Morales, 2018) ⁵¹	Other or mixed/Non- users	0. 8%	7	RR, 1.48 (1.29, 1.71)	22 877/ 2 400 720	6.8x10 ⁻⁸	24 (0.244)	1.09–2.02	No/NP	Yes	I	I / Moderate
Preterm birth (Huang, 2014) ⁶⁶	Other or mixed/Non- users	0. 4%	8	RR, 1.59 (1.31, 1.93)	3 506/ 910 029	3.4x10 ⁻⁷	35 (0.148)	1.02-2.47	No/NP	Yes	II	I / Moderate

Severe bleeding at any site (Laporte, 2017) ⁵⁵	Other or mixed/Non-users	6. 8%	44	OR, 1.41 (1.27,1.57)	190 016/ 1 512 411	2.2x10 ⁻¹⁰	90 (<0.000)	0.77-2.59	No/No	Yes	II	II / Low
Postpartum hemorrhage (Jiang, 2016) ⁵⁸	Other or mixed/Non-users	6. 8%	17	RR, 1.32 (1.17, 1.48)	49 155/ 65 1715	3.3x10 ⁻⁶	85 (<0.000)	0.84-2.07	No/NP	Yes	II	II / Low
Upper GI bleeding (Jiang, 2015) ⁶⁴	Other or mixed	0. 7%	22	OR, 1.55 (1.35, 1.78)	56 182/ 592 508	9.2x10 ⁻¹²	89 (<0.000)	0.83-2.91	No/No	Yes	II	II / Moderate
European studies												
Autism spectrum disorders (pregnancy maternal exposure; unadjusted estimates only; Andalib, 2017) ⁵²	SSRIs / Non-SSRIs users	0. 1%	4	OR, 1.80 (1.54, 2.10)	6 394/ 5 741 029	1.1x10 ⁻¹³	0 (0.396)	1.28–2.53	No/No	Yes	I	I/ Moderate
Osteoporotic fractures (Khanassov, 2018) ⁴²	SSRIs / Non-SSRIs users	6. 5%	12	RR, 1.76 (1.68, 1.87)	92 760/ 1 228 807	2.3x10 ⁻⁸⁶	67 (<0.000)	1.50-2.07	No/No	Yes	II	II/ Moderate
Upper GI bleeding (Jiang, 2015) ⁶⁴	SSRIs+others non-ADs/ No SSRIs use only+others non-ADs	0. 4%	14	OR, 1.60 (1.33, 1.93)	46 594/ 236 085	4.8x10 ⁻⁷	86 (<0.000)	0.80-3.91	No/NP	Yes	II	II / Moderate
Preterm birth (Huang, 2014) ⁶⁶	Any AD users/ No AD users	0. 9%	7	RR, 1.61 (1.36, 1.92)	17 518/ 1 984 543	5.6x10 ⁻⁸	56 (0.033)	1.02-2.55	No/NP	Yes	II	II / Moderate
Osteoporotic fractures (Wu, 2013) ⁷⁰	TCAs users/ Non-TCAs users	NA	6	RR, 1.37 (1.19, 1.56)	164 476/ 724 914	6.6x10 ⁻⁶	69 (0.007)	0.91-2.05	No/NP	Yes	II	II / Moderate

Hip Fracture (Oderda, 2012) ⁷⁶	TCAs and/or SSRIs users/ No AD users	2.1%	8	OR, 1.74 (1.39, 2.17)	40 196/ 159 706	1.1x10 ⁻⁶	89 (<0.000)	0.88-3.42	Yes/No	Yes	II	II / Critically low
North America studies												
Osteoporotic fractures (Khanassov, 2018) ⁴²	SSRIs / Non-SSRIs users	5.2 %	10	RR, 1.56 (1.33, 1.84)	31 683/ 249 808	1.0x10 ⁻⁷	87 (<0.000)	0.91-2.70	No/NP	Yes	II	II/ Moderate
Preterm birth (Huang, 2014) ⁶⁶	Any AD users/ No AD users	0.6%	17	RR, 1.70 (1.44, 1.99)	6 129/ 928 528	1.2x10 ⁻¹⁰	29 (0.124)	1.17-2.45	Yes/NP	Yes	II	II/ Moderate
Osteoporotic fractures (Wu, 2013) ⁷⁰	TCAs users/ Non-TCAs users	NA	6	RR, 1.54 (1.32, 1.81)	13 761 / 84 583	7.3 x10 ⁻⁸	76 (0.001)	0.95-2.51	No/No	Yes	II	II/ Moderate
Hip Fracture (Oderda, 2012) ⁷⁶	TCAs and/or SSRIs users/ No AD users	2.5%	8	OR, 1.81 (1.51, 2.18)	8 654/ 49 024	2.8x10 ⁻¹⁰	82 (<0.000)	1.01-3.27	No/NP	Yes	II	II/ Critically low
Other regions												
None	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA/ NA

AD–antidepressants, CE- class of evidence (main analysis), CES – class of evidence after sensitivity analysis, CI – confidence interval, ES – effect size, ESB –excess significance bias, GI–gastrointestinal, I²–heterogeneity, LS – largest study with significant effect, n – number of included studies per association, N – number of cases, NA– not applicable, NP – not pertinent because the number of observed studies is less than the expected, OR – odds ratio, PI – prediction interval, RR – relative risk, SNRIs – serotonin–norepinephrine reuptake inhibitors, SSRIs – selective serotonin reuptake inhibitors, SMD–standardized mean difference, SSE – small-study effect, TCAs – tricyclic antidepressants.

Supplement

eBox 1 Search strings for PubMed	2
eTable 1 Articles excluded after full-text revision, with reasons	5
eTable 2 Characteristics of meta-analyses of observational studies studying the association between antidepressants and risk of any adverse health outcome	11
eTable 3 Suggestive evidence (Class III) for the association of antidepressant use and risk of adverse health outcomes in meta-analyses of observational studies	16
eTable 4 Weak evidence (Class IV) for the association of antidepressant use and risk of adverse health outcomes in meta-analyses of observational studies.....	18
eTable 5 No evidence (Non-significant associations, $p>0.05$) for the association of antidepressant use and risk of adverse health outcomes in meta-analyses of observational studies.....	22
Supplementary methods.....	28
Selection between overlapping meta-analysis	28
Quality assessment	28
Grading method	29
Supplementary results	30
Sensitivity analysis results of convincing and highly suggestive evidence associations from the main analysis.....	30
<i>Cohort studies (both prospective and retrospective)</i>	<i>30</i>
<i>Prospective cohort studies</i>	<i>30</i>
<i>Studies adjusted for multiple covariates.....</i>	<i>30</i>
<i>Confounding by indication</i>	<i>31</i>
<i>High-quality primary studies</i>	<i>31</i>
<i>Classes of antidepressants</i>	<i>32</i>
<i>Studies located in Europe</i>	<i>32</i>
<i>Studies located in North America.....</i>	<i>33</i>
<i>Studies located in other regions.....</i>	<i>33</i>
Sensitivity analysis results of non-selected meta-analyses due to overlap.....	33
eTable 6 Sensitivity analysis results of non-selected meta-analyses due to overlap.....	35
eTable 7 List of covariates used for the sensitivity analysis limited to studies adjusted for covariates	42
PRISMA 2009 Checklist.....	50
MOOSE Checklist for Meta-analyses of Observational Studies	53
Supplementary references	55

eBox 1 Search strings for PubMed

("serotonin uptake inhibitors"[Pharmacological Action] OR "serotonin uptake inhibitors"[MeSH Terms] OR ("serotonin"[All Fields] AND "uptake"[All Fields] AND "inhibitors"[All Fields]) OR "serotonin uptake inhibitors"[All Fields] OR ("selective"[All Fields] AND "serotonin"[All Fields] AND "reuptake"[All Fields] AND "inhibitors"[All Fields]) OR "selective serotonin reuptake inhibitors"[All Fields]) OR ("serotonin uptake inhibitors"[Pharmacological Action] OR "serotonin uptake inhibitors"[MeSH Terms] OR ("serotonin"[All Fields] AND "uptake"[All Fields] AND "inhibitors"[All Fields]) OR "serotonin uptake inhibitors"[All Fields] OR "ssris"[All Fields]) OR ("antidepressive agents"[Pharmacological Action] OR "antidepressive agents"[MeSH Terms] OR ("antidepressive"[All Fields] AND "agents"[All Fields]) OR "antidepressive agents"[All Fields] OR "antidepressants"[All Fields]) OR (antidepranalgesia[All Fields] OR antidepredatory[All Fields] OR antideprein[All Fields] OR antidepres[All Fields] OR antidepres'iva[All Fields] OR antidepres'ivn'i[All Fields] OR antidepres'ivn'ich[All Fields] OR antidepres'ivy[All Fields] OR antidepresan[All Fields] OR antidepresant[All Fields] OR antidepresanti[All Fields] OR antidepresants[All Fields] OR antidepresantu[All Fields] OR antidepresessant[All Fields] OR antidepreseurs[All Fields] OR antidepresiv[All Fields] OR antidepresiva[All Fields] OR antidepresivach[All Fields] OR antidepresivami[All Fields] OR antidepresivas[All Fields] OR antidepressive[All Fields] OR antidepresivem[All Fields] OR antidepressives[All Fields] OR antidepressivi[All Fields] OR antidepressivima[All Fields] OR antidepressivnm[All Fields] OR antidepressivnaih[All Fields] OR antidepressivni[All Fields] OR antidepressivnich[All Fields] OR antidepressivniho[All Fields] OR antidepressivnim[All Fields] OR antidepressivnimu[All Fields] OR antidepressivnych[All Fields] OR antidepressivo[All Fields] OR antidepressivos[All Fields] OR antidepressivum[All Fields] OR antidepressivy[All Fields] OR antidepresor[All Fields] OR antidepress[All Fields] OR antidepressand[All Fields] OR antidepressani[All Fields] OR antidepressanov[All Fields] OR antidepressans[All Fields] OR antidepressant[All Fields] OR antidepressant'[All Fields] OR antidepressant's[All Fields] OR antidepressanta[All Fields] OR antidepressantam[All Fields] OR antidepressantami[All Fields] OR antidepressanteffects[All Fields] OR antidepressantinduced[All Fields] OR antidepressantlike[All Fields] OR antidepressantlithium[All Fields] OR antidepressantnogo[All Fields] OR antidepressantom[All Fields] OR antidepressantov[All Fields] OR antidepressants[All Fields] OR antidepressants'[All Fields] OR antidepressantswere[All Fields] OR antidepressantu[All Fields] OR antidepressantv[All Fields] OR antidepressanty[All Fields] OR antidepressed[All Fields] OR antidepressent[All Fields] OR antidepressents[All Fields] OR antidepresseur[All Fields] OR antidepresseurs[All Fields] OR antidepresseurs'[All Fields] OR antidepresseus[All Fields] OR antidepresseuses[All Fields] OR antidepressia[All Fields] OR antidepressief[All Fields] OR antidepressieve[All Fields] OR antidepressiewe[All Fields] OR antidepressiff[All Fields] OR antidepressifs[All Fields] OR antidepressiivien[All Fields] OR antidepressiiviset[All Fields] OR antidepressin[All Fields] OR antidepressing[All Fields] OR antidepression[All Fields] OR antidepressions[All Fields] OR antidepressionsbehandling[All Fields] OR antidepressiu[All Fields] OR antidepressiv[All Fields] OR antidepressiva[All Fields] OR antidepressivabehandling[All Fields] OR antidepressivarichtlijnen[All Fields] OR antidepressivas[All Fields] OR antidepressivastudien[All Fields] OR antidepressivasubstanzen[All Fields] OR antidepressivatherapie[All Fields] OR antidepressive[All Fields] OR antidepressively[All Fields] OR antidepressivem[All Fields] OR antidepressiven[All Fields] OR antidepressivene[All Fields] OR antidepressiver[All Fields] OR antidepressives[All Fields] OR antidepressivi[All Fields] OR antidepressivnikh[All Fields] OR antidepressivnoe[All Fields] OR antidepressivnogo[All Fields] OR antidepressivnoi[All Fields] OR antidepressivnom[All Fields] OR antidepressivnye[All Fields] OR antidepressivnyi[All Fields] OR antidepressivnykh[All Fields] OR antidepressivnymi[All Fields] OR antidepressivo[All Fields] OR antidepressivos[All Fields] OR antidepressivt[All Fields] OR antidepressivum[All Fields] OR antidepressivum'[All Fields] OR antidepressivums[All Fields] OR

antidepressziv[All Fields] OR antidepresszivum[All Fields] OR antidepresszivumhasznalat[All Fields] OR antidepresszivumok[All Fields] OR antidepresszivumot[All Fields] OR antideprimenti[All Fields] OR antidepressin[All Fields] OR antidepresssants[All Fields] OR antidepresssiva[All Fields] OR antidepresssives[All Fields]) OR ("citalopram"[MeSH Terms] OR "citalopram"[All Fields]) OR ("citalopram"[MeSH Terms] OR "citalopram"[All Fields] OR "escitalopram"[All Fields]) OR ("fluoxetine"[MeSH Terms] OR "fluoxetine"[All Fields]) OR ("fluvoxamine"[MeSH Terms] OR "fluvoxamine"[All Fields]) OR ("paroxetine"[MeSH Terms] OR "paroxetine"[All Fields]) OR ("sertraline"[MeSH Terms] OR "sertraline"[All Fields]) OR ("monoamine oxidase inhibitors"[Pharmacological Action] OR "monoamine oxidase inhibitors"[MeSH Terms] OR ("monoamine"[All Fields] AND "oxidase"[All Fields] AND "inhibitors"[All Fields]) OR "monoamine oxidase inhibitors"[All Fields] OR "maoi"[All Fields]) OR ("antidepressive agents, tricyclic"[Pharmacological Action] OR "antidepressive agents, tricyclic"[MeSH Terms] OR ("antidepressive"[All Fields] AND "agents"[All Fields] AND "tricyclic"[All Fields]) OR "tricyclic antidepressive agents"[All Fields] OR ("tricyclic"[All Fields] AND "antidepressant"[All Fields]) OR "tricyclic antidepressant"[All Fields]) OR ("Theor Chem Acc"[Journal] OR "tca"[All Fields]) OR (("serotonin"[MeSH Terms] OR "serotonin"[All Fields]) AND ("norepinephrine"[MeSH Terms] OR "norepinephrine"[All Fields] OR "noradrenaline"[All Fields]) AND reuptake[All Fields] AND inhibitor[All Fields]) OR ("serotonin and noradrenaline reuptake inhibitors"[Pharmacological Action] OR "serotonin and noradrenaline reuptake inhibitors"[MeSH Terms] OR ("serotonin"[All Fields] AND "noradrenaline"[All Fields] AND "reuptake"[All Fields] AND "inhibitors"[All Fields]) OR "serotonin and noradrenaline reuptake inhibitors"[All Fields] OR "snri"[All Fields]) OR (("serotonin antagonists"[Pharmacological Action] OR "serotonin antagonists"[MeSH Terms] OR ("serotonin"[All Fields] AND "antagonists"[All Fields]) OR "serotonin antagonists"[All Fields] OR ("serotonin"[All Fields] AND "antagonist"[All Fields]) OR "serotonin antagonist"[All Fields]) AND reuptake[All Fields] AND inhibitor[All Fields]) OR SARI[All Fields] OR (("norepinephrine"[MeSH Terms] OR "norepinephrine"[All Fields]) AND ("dopamine uptake inhibitors"[Pharmacological Action] OR "dopamine uptake inhibitors"[MeSH Terms] OR ("dopamine"[All Fields] AND "uptake"[All Fields] AND "inhibitors"[All Fields]) OR "dopamine uptake inhibitors"[All Fields] OR ("dopamine"[All Fields] AND "reuptake"[All Fields] AND "inhibitor"[All Fields]) OR "dopamine reuptake inhibitor"[All Fields])) OR NDRI[All Fields] OR (("norepinephrine"[MeSH Terms] OR "norepinephrine"[All Fields]) AND reuptake[All Fields] AND inhibitor[All Fields]) OR NRI[All Fields] OR (noradrenergic[All Fields] AND specific[All Fields] AND serotonergic[All Fields] AND ("antidepressive agents"[Pharmacological Action] OR "antidepressive agents"[MeSH Terms] OR ("antidepressive"[All Fields] AND "agents"[All Fields]) OR "antidepressive agents"[All Fields] OR "antidepressant"[All Fields])) OR NaSSA[All Fields] AND (("suicide"[MeSH Terms] OR "suicide"[All Fields]) OR ("suicidal ideation"[MeSH Terms] OR ("suicidal"[All Fields] AND "ideation"[All Fields]) OR "suicidal ideation"[All Fields]) OR ("suicide"[MeSH Terms] OR "suicide"[All Fields]) AND ("risk"[MeSH Terms] OR "risk"[All Fields])) OR (serious[All Fields] AND adverse[All Fields] AND events[All Fields]) OR ("adverse effects"[Subheading] OR ("adverse"[All Fields] AND "effects"[All Fields]) OR "adverse effects"[All Fields] OR ("side"[All Fields] AND "effects"[All Fields]) OR "side effects"[All Fields]) OR harm[All Fields] OR ("fractures, bone"[MeSH Terms] OR ("fractures"[All Fields] AND "bone"[All Fields]) OR "bone fractures"[All Fields] OR ("bone"[All Fields] AND "fractures"[All Fields])) OR ("psychomotor agitation"[MeSH Terms] OR ("psychomotor"[All Fields] AND "agitation"[All Fields]) OR "psychomotor agitation"[All Fields] OR "akathisia"[All Fields]) OR ("photosensitivity disorders"[MeSH Terms] OR ("photosensitivity"[All Fields] AND "disorders"[All Fields]) OR "photosensitivity disorders"[All Fields] OR "photosensitivity"[All Fields]) OR ("sexual behavior"[MeSH Terms] OR ("sexual"[All Fields] AND "behavior"[All Fields]) OR "sexual behavior"[All Fields] OR "sexual"[All Fields]) AND ("physiopathology"[Subheading] OR "physiopathology"[All Fields] OR "dysfunction"[All Fields])) OR ("coronary disease"[MeSH Terms] OR ("coronary"[All Fields] AND "disease"[All Fields]) OR "coronary disease"[All Fields] OR ("coronary"[All Fields] AND "heart"[All Fields] AND "disease"[All Fields]) OR

"coronary heart disease"[All Fields]) OR ("hemorrhage"[MeSH Terms] OR "hemorrhage"[All Fields] OR "bleeding"[All Fields]) OR (("fractures, bone"[MeSH Terms] OR ("fractures"[All Fields] AND "bone"[All Fields]) OR "bone fractures"[All Fields] OR "fracture"[All Fields]) AND ("risk"[MeSH Terms] OR "risk"[All Fields])) OR (discontinuation[All Fields] AND ("syndrome"[MeSH Terms] OR "syndrome"[All Fields])) OR ("serotonin syndrome"[MeSH Terms] OR ("serotonin"[All Fields] AND "syndrome"[All Fields]) OR "serotonin syndrome"[All Fields]) OR ("neonatal abstinence syndrome"[MeSH Terms] OR ("neonatal"[All Fields] AND "abstinence"[All Fields] AND "syndrome"[All Fields]) OR "neonatal abstinence syndrome"[All Fields]) OR ("autistic disorder"[MeSH Terms] OR ("autistic"[All Fields] AND "disorder"[All Fields]) OR "autistic disorder"[All Fields] OR "autism"[All Fields])) AND (("meta-analysis"[Publication Type] OR "meta-analysis as topic"[MeSH Terms] OR "meta-analysis"[All Fields]) OR ("meta-analysis as topic"[MeSH Terms] OR ("meta-analysis"[All Fields] AND "topic"[All Fields]) OR "meta-analysis as topic"[All Fields] OR "metaanalysis"[All Fields]) OR "systematic review"[All Fields])

eTable 1 Articles excluded after full-text revision, with reasons

Author, year (Reference)	Reason for exclusion
Ng, 2019 ¹	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Patel, 2019 ²	Not reporting an association of antidepressants and risk of adverse health outcomes
Rexwinkel, 2019 ³	Nether a meta-analysis or systematic review with quantitative synthesis
Perlman, 2019 ⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Amare, 2019 ⁵	Not reporting an association of antidepressants and risk of adverse health outcomes
Aguiar, 2019 ⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Zhou, 2018 ⁷	Not included only case-control and cohort studies
Wu, 2018 ⁸	Not reporting an association of antidepressants and risk of adverse health outcomes
Wang, 2018 ⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Visco, 2018 ¹⁰	Not included only case-control and cohort studies
Uguz, 2018 ¹¹	Nether a meta-analysis or systematic review with quantitative synthesis
Tan, 2018 ¹²	Not a meta-analysis or systematic review of observational studies
Telang, 2018 ¹³	Not a meta-analysis or systematic review of observational studies
Sepehrpour, 2018 ¹⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Prady, 2018 ¹⁵	Nether a meta-analysis or systematic review with quantitative synthesis
Cappetta, 2018 ¹⁶	Not a meta-analysis or systematic review of observational studies
Steinert, 2018 ¹⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Melo, 2018 ¹⁸	Not included only case-control and cohort studies
Moncrieff, 2018 ¹⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Douros, 2018 ²⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Chen, 2018 ²¹	Not a meta-analysis or systematic review of observational studies
Comoretto, 2018 ²²	Not reporting an association of antidepressants and risk of adverse health outcomes
Varney, 2017 ²³	Nether a meta-analysis or systematic review with quantitative synthesis
Uguz, 2017 ²⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Salvi, 2017 ²⁵	Not included only case-control and cohort studies
Stubbs, 2017 ²⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Riediger, 2017 ²⁷	Not a meta-analysis or systematic review of observational studies
Querido, 2017 ²⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Mezzacappa, 2017 ²⁹	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Maund, 2017 ³⁰	Not a meta-analysis or systematic review of observational studies
Moraros, 2017 ³¹	Not included only case-control and cohort studies
Locher, 2017 ³²	Not a meta-analysis or systematic review of observational studies
Laux, 2017 ³³	Nether a meta-analysis or systematic review with quantitative synthesis
Kaplan, 2017 ³⁴	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Heller, 2017 ³⁵	Nether a meta-analysis or systematic review with quantitative synthesis
Hill, 2017 ³⁶	Not included only case-control and cohort studies
Deidda, 2017 ³⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Brunnauer, 2017 ³⁸	Nether a meta-analysis or systematic review with quantitative synthesis

Brown, 2017 ³⁹	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Beyer, 2017 ⁴⁰	Not a meta-analysis or systematic review of observational studies
Andrade, 2017 ⁴¹	Nether a meta-analysis or systematic review with quantitative synthesis
Allain, 2017 ⁴²	Nether a meta-analysis or systematic review with quantitative synthesis
Vlachos, 2016 ⁴³	Nether a meta-analysis or systematic review with quantitative synthesis
Werneke, 2016 ⁴⁴	Not reporting an association of antidepressants and risk of adverse health outcomes
Warden, 2016 ⁴⁵	Nether a meta-analysis or systematic review with quantitative synthesis
Torjesen, 2016 ⁴⁶	Commentary
Stone, 2016 ⁴⁷	Commentary
Smit, 2016 ⁴⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Simonsen, 2016 ⁴⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Sharma, 2016 ⁵⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Selmer, 2016 ⁵¹	Insufficient or inadequate data for quantitative synthesis provided
Rudisill, 2016 ⁵²	Insufficient or inadequate data for quantitative synthesis provided
Pozzi, 2016 ⁵³	Nether a meta-analysis or systematic review with quantitative synthesis
Potočnjak, 2016 ⁵⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Muzik, 2016 ⁵⁵	Nether a meta-analysis or systematic review with quantitative synthesis
Lassen, 2016 ⁵⁶	Insufficient or inadequate data for quantitative synthesis provided
Kaplan, 2016 ⁵⁷	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Kobayashi, 2016 ⁵⁸	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Jordan, 2016 ⁵⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Jarde, 2016 ⁶⁰	Not reporting an association of antidepressants and risk of adverse health outcomes
Howland, 2016 ⁶¹	Nether a meta-analysis or systematic review with quantitative synthesis
Galling, 2016 ⁶²	Not reporting an association of antidepressants and risk of adverse health outcomes
Donneyong, 2016 ⁶³	Nether a meta-analysis or systematic review with quantitative synthesis
Cameron, 2016 ⁶⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Braun, 2016 ⁶⁵	Not a meta-analysis or systematic review of observational studies
Bielefeldt, 2016 ⁶⁶	Not a meta-analysis or systematic review of observational studies
Barth, 2016 ⁶⁷	Not a meta-analysis or systematic review of observational studies
Akiyamen, 2016 ⁶⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Alvares, 2016 ⁶⁹	Not included only case-control and cohort studies
Wang, 2015 ⁷⁰	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Uchida, 2015 ⁷¹	Nether a meta-analysis or systematic review with quantitative synthesis
Thase, 2015 ⁷²	Nether a meta-analysis or systematic review with quantitative synthesis
Teo, 2015 ⁷³	Nether a meta-analysis or systematic review with quantitative synthesis
Tampi, 2015 ⁷⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Stubbs, 2015 ⁷⁵	Commentary
Santarsieri, 2015 ⁷⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Robinson, 2015 ⁷⁷	Commentary
Renoux, 2015 ⁷⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Pinheiro, 2015 ⁷⁹	Insufficient or inadequate data for quantitative synthesis provided

Orsolini, 2015 ⁸⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Olfson, 2015 ⁸¹	Nether a meta-analysis or systematic review with quantitative synthesis
Nezafati, 2015 ⁸²	Nether a meta-analysis or systematic review with quantitative synthesis
Mohler, 2015 ⁸³	Commentary
Man, 2015 ⁸⁴	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
KoKoAung, 2015 ⁸⁵	Full-text could not retrieved
Götzsche, 2015 ⁸⁶	Commentary
Gentile, 2015 ⁸⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Gentile, 2015 ⁸⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Gebara, 2015 ⁸⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Ennis, 2015 ⁹⁰	Not reporting an association of antidepressants and risk of adverse health outcomes
Davis, 2015 ⁹¹	Not reporting an association of antidepressants and risk of adverse health outcomes
Correll, 2015 ⁹²	Nether a meta-analysis or systematic review with quantitative synthesis
Bruning, 2015 ⁹³	Nether a meta-analysis or systematic review with quantitative synthesis
Stevenson, 2014 ⁹⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Ross, 2014 ⁹⁵	Commentary
Rais, 2014 ⁹⁶	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Park, 2015 ⁹⁷	Not included only case-control and cohort studies
Prieto-Alhambra, 2014 ⁹⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Previti, 2014 ⁹⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Pereira, 2014 ¹⁰⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Pedersen, 2014 ¹⁰¹	Nether a meta-analysis or systematic review with quantitative synthesis
Paumgartten, 2014 ¹⁰²	Nether a meta-analysis or systematic review with quantitative synthesis
Okazaki, 2014 ¹⁰³	Nether a meta-analysis or systematic review with quantitative synthesis
Oka, 2014 ¹⁰⁴	Insufficient or inadequate data for quantitative synthesis provided
McDonagh, 2014 ¹⁰⁵	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Mahdanian, 2014 ¹⁰⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Mago, 2014 ¹⁰⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Lorenzo, 2014 ¹⁰⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Li, 2014 ¹⁰⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Isacsson, 2014 ¹¹⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Grigoriadis, 2014 ¹¹¹	Nether a meta-analysis or systematic review with quantitative synthesis
Gebara, 2014 ¹¹²	Nether a meta-analysis or systematic review with quantitative synthesis
Fanoe, 2014 ¹¹³	Nether a meta-analysis or systematic review with quantitative synthesis
El Marroun, 2014 ¹¹⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Costoloni, 2014 ¹¹⁵	Nether a meta-analysis or systematic review with quantitative synthesis
Clayton, 2014 ¹¹⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Beach, 2014 ¹¹⁷	Insufficient or inadequate data for quantitative synthesis provided
Anglin, 2014 ¹¹⁸	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Andrade, 2014 ¹¹⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Rotella, 2013 ¹²⁰	Not reporting an association of antidepressants and risk of adverse health outcomes

Rabenda, 2013 ¹²¹	Insufficient or inadequate data for quantitative synthesis provided
Painuly, 2013 ¹²²	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Nassir, 2013 ¹²³	Nether a meta-analysis or systematic review with quantitative synthesis
Myles, 2013 ¹²⁴	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Mintzes, 2013 ¹²⁵	Commentary
Kennedy, 2013 ¹²⁶	Commentary
Howland, 2013 ¹²⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Grigoriadis, 2013 ¹²⁸	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Grigoriadis, 2013 ¹²⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Gahr, 2013 ¹³⁰	Nether a meta-analysis or systematic review with quantitative synthesis
De Jong, 2013 ¹³¹	Nether a meta-analysis or systematic review with quantitative synthesis
De Groot, 2013 ¹³²	Nether a meta-analysis or systematic review with quantitative synthesis
Brunnauer, 2013 ¹³³	Nether a meta-analysis or systematic review with quantitative synthesis
Bhattacharjee, 2013 ¹³⁴	Not included only case-control and cohort studies
Barnard, 2013 ¹³⁵	Nether a meta-analysis or systematic review with quantitative synthesis
Al-Zoairy, 2013 ¹³⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Wu, 2012 ¹³⁷	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
T Jong, 2012 ¹³⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Sansone, 2012 ¹³⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Pan, 2012 ¹⁴⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Oyebode, 2012 ¹⁴¹	Nether a meta-analysis or systematic review with quantitative synthesis
Nischal, 2012 ¹⁴²	Nether a meta-analysis or systematic review with quantitative synthesis
Malm, 2012 ¹⁴³	Nether a meta-analysis or systematic review with quantitative synthesis
Lopez-Yarto, 2012 ¹⁴⁴	Insufficient or inadequate data for quantitative synthesis provided
Kucukaycan, 2012 ¹⁴⁵	Nether a meta-analysis or systematic review with quantitative synthesis
KoKoAung, 2012 ¹⁴⁶	Full-text could not retrieved
Hennings, 2012 ¹⁴⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Hackam, 2012 ¹⁴⁸	Insufficient or inadequate data for quantitative synthesis provided
Grzeskowiak, 2012 ¹⁴⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Einarson, 2012 ¹⁵⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Eom, 2012 ¹⁵¹	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Colotto, 2012 ¹⁵²	Nether a meta-analysis or systematic review with quantitative synthesis
Bromley, 2012 ¹⁵³	Nether a meta-analysis or systematic review with quantitative synthesis
Adams, 2012 ¹⁵⁴	Not reporting an association of antidepressants and risk of adverse health outcomes
Gentile, 2011 ¹⁵⁵	Nether a meta-analysis or systematic review with quantitative synthesis
Gentile, 2011 ¹⁵⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Fenger-Gron, 2011 ¹⁵⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Davanzo, 2011 ¹⁵⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Dassanayake, 2011 ¹⁵⁹	Not reporting an association of antidepressants and risk of adverse health outcomes
Berkowitz, 2011 ¹⁶⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Barbui, 2011 ¹⁶¹	Nether a meta-analysis or systematic review with quantitative synthesis

Wurst, 2010 ¹⁶²	Insufficient or inadequate data for quantitative synthesis provided
Wu, 2010 ¹⁶³	Not reporting an association of antidepressants and risk of adverse health outcomes
Van Driel, 2010 ¹⁶⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Udechuku, 2010 ¹⁶⁵	Nether a meta-analysis or systematic review with quantitative synthesis
Tuccori, 2010 ¹⁶⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Simoncelli, 2010 ¹⁶⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Scialli, 2010 ¹⁶⁸	Commentary
Kemp, 2010 ¹⁶⁹	Not included only case-control and cohort studies
Kölch, 2010 ¹⁷⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Kontakioti, 2010 ¹⁷¹	Nether a meta-analysis or systematic review with quantitative synthesis
Einarson, 2010 ¹⁷²	Commentary
Einarson, 2010 ¹⁷³	Nether a meta-analysis or systematic review with quantitative synthesis
Berard, 2010 ¹⁷⁴	Commentary
Woolcott, 2009 ¹⁷⁵	Not included only case-control and cohort studies
Fortinguerra, 2009 ¹⁷⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Bergemann, 2009 ¹⁷⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Taylor, 2008 ¹⁷⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Sterke, 2008 ¹⁷⁹	Nether a meta-analysis or systematic review with quantitative synthesis
O'Brien, 2008 ¹⁸⁰	Insufficient or inadequate data for quantitative synthesis provided
Loke, 2008 ¹⁸¹	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Kongkaew, 2008 ¹⁸²	Nether a meta-analysis or systematic review with quantitative synthesis
Bond, 2008 ¹⁸³	Not a meta-analysis or systematic review of observational studies
Takkouche, 2007 ¹⁸⁴	Insufficient or inadequate data for quantitative synthesis provided
Howard, 2007 ¹⁸⁵	Nether a meta-analysis or systematic review with quantitative synthesis
Hartikainen, 2007 ¹⁸⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Bellantuono, 2007 ¹⁸⁷	Nether a meta-analysis or systematic review with quantitative synthesis
Bar-Oz, 2007 ¹⁸⁸	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
Swenson, 2006 ¹⁸⁹	Not a meta-analysis or systematic review of observational studies
Sala, 2006 ¹⁹⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Rahimi, 2006 ¹⁹¹	Overlapping meta-analysis or systematic review of association of antidepressants and risk of adverse health outcomes
McIntyre, 2006 ¹⁹²	Nether a meta-analysis or systematic review with quantitative synthesis
McClintock, 2006 ¹⁹³	Nether a meta-analysis or systematic review with quantitative synthesis
Hall, 2006 ¹⁹⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Lattimore, 2005 ¹⁹⁵	Insufficient or inadequate data for quantitative synthesis provided
Gentile, 2005 ¹⁹⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Einarson, 2005 ¹⁹⁷	Insufficient or inadequate data for quantitative synthesis provided
Wen, 2004 ¹⁹⁸	Nether a meta-analysis or systematic review with quantitative synthesis
Weissman, 2004 ¹⁹⁹	Nether a meta-analysis or systematic review with quantitative synthesis
Ramasubbu, 2004 ²⁰⁰	Nether a meta-analysis or systematic review with quantitative synthesis
Newman, 2004 ²⁰¹	Nether a meta-analysis or systematic review with quantitative synthesis
McIntosh, 2004 ²⁰²	Nether a meta-analysis or systematic review with quantitative synthesis

Bailey, 2004 ²⁰³	Insufficient or inadequate data for quantitative synthesis provided
Weinrieb, 2003 ²⁰⁴	Nether a meta-analysis or systematic review with quantitative synthesis
Addis, 2000 ²⁰⁵	Insufficient or inadequate data for quantitative synthesis provided
Thase, 1998 ²⁰⁶	Nether a meta-analysis or systematic review with quantitative synthesis
Wilens, 1996 ²⁰⁷	Nether a meta-analysis or systematic review with quantitative synthesis

eTable 2 Characteristics of meta-analyses of observational studies studying the association between antidepressants and risk of any adverse health outcome

Study (see references 38-81 in the main text)	Type of studies included	Risks of adverse health outcomes examined	Exposures	Non-exposures (comparator)	No of associations (no of included studies estimates)	Population (s)	Average of adjustments (range)	AMSTAR2
Masarwa, 2019	Case-control and cohort	Persistent pulmonary hypertension of the newborn	SSRIs+SNRIs	No SSRIs+SNRIs use	2 (19)	Pregnant women (any trimester)	9 (4-16)	Moderate
Halvorsen, 2019	Case-control and cohort	Autism spectrum disorders, attention-deficit hyperactivity disorder and mental retardation in children	SSRIs	No SSRIs use	3 (16)	Pregnant women (any trimester)	12 (4-19)	Moderate
Wang, 2018	Case-control and cohort	Dementia	SSRIs; TCAs; MAOIs	No ADs use	3 (11)	Elderly patients with various disorders	10 (8-14)	Moderate
Schweiger, 2018	Case-control and cohort	Bone mineral density	SSRIs; TCAs	No ADs use	4 (8)	Women with depressive disorders	NR	Critically Low
Khanassov, 2018	Case-control and cohort	Osteoporotic fractures	SSRIs	No SSRIs	1 (24)	General population of adults and older adults	15 (2-29)	Moderate
Jiang, 2018	Cohort	Attention-deficit hyperactivity disorder in children	Any AD	No ADs use	8 (34)	Pregnant women (any trimester)	9 (7-13)	Moderate
Gao, 2018	Cohort	Congenital malformations in infants	SSRIs	No SSRIs use	7 (37)	Pregnant women (first trimester)	5 (2-9)	Moderate
Guan, 2018	Case-control and cohort	Gestational hypertension and/or preeclampsia	SSRIs	No SSRIs use	3 (16)	Pregnant women (any trimester)	14 (10-18)	Moderate
Chappuis, 2018	Cohort	Dental implant failure	SSRIs	No SSRIs use	1 (2)	General adult population	4 (2-6)	Moderate
Na, 2018	Case-control	GI bleeding; any other type of bleeding; both GI and any other type of bleeding	Mirtazapine; Bupropion	SSRIs; No medication	5 (20)	Patients with a diagnosis of abnormal bleeding; warfarin	9 (4-16)	Moderate

Study (see references 38-81 in the main text)	Type of studies included	Risks of adverse health outcomes examined	Exposures	Non-exposures (comparator)	No of associations (no of included studies estimates)	Population (s)	Average of adjustments (range)	AMSTAR2
						users; psychiatric inpatients		
Man, 2018	Case-control and cohort	Attention-deficit hyperactivity disorder in children	Any AD	No ADs use	2 (12)	Women with prenatal or pre-conception exposure to antidepressants	8 (7-9)	Moderate
Fu, 2018	Nested case-control and case-control	Cataract risk	SSRIs; SNRIs; TCAs	No ADs use	3 (13)	Patients with medical/psychiatric diagnoses	5 (3-10)	Moderate
Eckersley, 2018	Cohort	Bleeding; mortality	SSRIs	No SSRIs use	4 (14)	Patients undergoing coronary artery bypass graft surgery	13 (9-16)	Moderate
Zhou, 2018	Case-control and cohort	Autism spectrum disorders	Any AD	No ADs use	1 (13)	Pregnant women (any trimester)	8 (3-11)	Low
Morales, 2018	Case-control and cohort	Autism spectrum disorders	Any AD	No ADs use	2 (13)	Women with pre-conception exposure to antidepressants	8 (3-11)	Moderate
Andalib, 2017	Case-control and cohort	Autism spectrum disorders	SSRIs	No SSRIs use	1 (7)	Pregnant women (any trimester)	7 (3-11)	Moderate
Zhang, 2017	Cohort	Cardiovascular-related malformations of infants; Both atrial and ventricular septal defect	SSRIs	No SSRIs use	2 (37)	Pregnant women (first trimester)	6 (0-13)	Low
Shen, 2017	Cohort	Cardiovascular-related malformations of infants	Sertraline	No sertraline or any other AD use	6 (40)	Pregnant women (first trimester)	6 (0-9)	Moderate
Laporte, 2017	Case-control and cohort	Severe bleeding at any site	SSRIs; S NRIs	No AD; No SSRI; Other AD; Neither SSRI nor NSAID	1 (44)	Patients with a diagnosis of abnormal	6 (0-17)	Low

Study (see references 38-81 in the main text)	Type of studies included	Risks of adverse health outcomes examined	Exposures	Non-exposures (comparator)	No of associations (no of included studies estimates)	Population (s)	Average of adjustments (range)	AMSTAR2
						bleeding; surgical patients		
Gao, 2017	Cohort	Congenital malformations in infants	Fluoxetine	No ADs or teratogens use	8 (38)	Pregnant women (first trimester)	5 (2-9)	Moderate
Biffi, 2017	Case-control and cohort	Onset of acute heart disease; Cerebrovascular disease	SSRIs; TCAs; Any other AD	No ADs use	4 (31)	Patients with depression	9 (0-16)	Moderate
Jiang, 2016	Nested case-control, case-control and cohort	Postpartum hemorrhage	SSRIs; Any other non-SSRIs use	No ADs use	1 (17)	Pregnant women (any trimester)	13 (7-22)	Low
Healy, 2016	Cohort	Neurodevelopmental delay/spectrum disorders	SSRI	No SSRIs	1 (17)	Pregnant women (any trimester)	NR	Moderate
Eke, 2016	Cohort	Preterm birth; respiratory distress syndrome	SSRIs	No SSRIs	3 (16)	Pregnant women (first and third trimester)	1.5 (0-5)	Moderate
Bérard, 2016	Case-control and cohort	Major cardiac malformations	Paroxetine	No paroxetine use or no any other ADs use	6 (87)	Pregnant women (first trimester)	6 (1-15)	Moderate
Undela, 2015	Case-control and cohort	Myocardial infarction	TCAs; SSRIs; Any other AD	No ADs use	1 (21)	Patients with depression	10 (2-18)	Moderate
Singh, 2015	Cohort	Perioperative bleeding outcomes and/or any cause mortality	SSRIs+SNRIs; Any other ADs	No SSRIs+SNRIs use or no any other ADs use	4 (19)	Pre-operative patients	8.6 (1-16)	Low
Jiang, 2015	Case-control and cohort	Upper GI bleeding	SSRIs+ NSAIDs + acid-suppressing drugs+ antiplatelet drugs	No ADs use only	2 (30)	Patients with various disorders with a diagnosis of UGIB	9 (0-17)	Moderate

Study (see references 38-81 in the main text)	Type of studies included	Risks of adverse health outcomes examined	Exposures	Non-exposures (comparator)	No of associations (no of included studies estimates)	Population (s)	Average of adjustments (range)	AMSTAR2
Shin, 2014	Case-control and cohort	Stroke (intracerebral hemorrhage and subarachnoid hemorrhage)	SSRIs	No SSRIs	3 (23)	Patients with depression	13 (4-31)	Low
Huang, 2014	Case-control and cohort	Low birth weight; Preterm birth	SSRIs; Any other non-SSRIs use	No ADs use	2 (43)	Pregnant women (any trimester)	NR	Moderate
Huybrechts, 2014	Case-control and cohort	Spontaneous abortion	SSIRs	No SSRIs	2 (20)	Pregnant women (early and late pregnancy)	NR	Moderate
Grigoriadis, 2014	Case-control and cohort	Pulmonary hypertension of the newborn	SSRIs	No SSRIs or no any other ADs use	1 (3)	Pregnant women with (early pregnancy)	7 (4-10)	Moderate
Oh, 2014	Case-control and cohort	Coronary heart disease	TCAs	No TCAs	1 (14)	General population of adults	NR	Moderate
Wu, 2013	Case-control and cohort	Osteoporotic fractures	TCAs	No TCAs	1 (12)	General population of older adults	12 (2-30)	Moderate
Ross, 2013	Cohort	Gestational age; APGAR score	TCAs; SSRIs; Any other AD	No ADs use	3 (42)	Pregnant women (any trimester)	3 (0-21)	Moderate
Riggin, 2013	Cohort	Congenital malformations in infants	Fluoxetine	No fluoxetine or teratogens use	2 (38)	Pregnant women (first trimester)	NR	Moderate
Grigoriadis, 2013	Case-control and cohort	Poor neonatal adaptation; Respiratory distress syndrome; Tremors	TCAs; SSRIs; SNRIs;Any other AD	No ADs use	3 (21)	Pregnant women (any trimester)	5 (4-7)	Moderate
Myles, 2013	Case-control and cohort	Major malformations in infants	Citalopram	No ADs use	2 (13)	Pregnant women (first trimester)	NR	High quality
Nikfar, 2012	Case-control and cohort	Major malformations in infants; Spontaneous abortion	SSRIs	No SSRIs	2 (28)	Pregnant women (any trimester)	NR	Low
Oderda, 2012	Case-control and cohort	Hip Fracture	Any AD	No ADs use	1 (18)	General population of older adults	5 (0-19)	Critically Low
Eom, 2012	Case-control and cohort	Breast cancer	Any AD	No ADs use	1 (18)	General population of adults	7 (1-13)	Moderate

Study (see references 38-81 in the main text)	Type of studies included	Risks of adverse health outcomes examined	Exposures	Non-exposures (comparator)	No of associations (no of included studies estimates)	Population (s)	Average of adjustments (range)	AMSTAR2
Lee, 2012	Case-control and cohort	Colon cancer	SSRIs	No SSRIs or no any other ADs use	1 (6)	General population of adults and older adults	6 (2-7)	Moderate
Cosgrove, 2011	Case-control and cohort	Breast and ovarian cancer	SSRIs; TCAs	No ADs use	2 (32)	General population of adults	NR	Low
Barbui, 2009	Case-control and cohort	Suicide attempt and completion	SSRIs	No SSRIs	3 (14)	Patients with major depressive disorders	6 (4-8)	High quality
Hemels, 2005	Cohort	Spontaneous abortion	Any AD	No ADs use	1 (11)	Pregnant women (any trimester)	NR	Moderate

AD-antidepressants, AMSTAR2 – acronym of A Measurement Tool to Assess systematic Reviews, version 2, BMD – bone mineral density, APGAR – acronym of Appearance (skin color), Pulse (heart rate), Grimace (reflex irritability), Activity (muscle tone), and Respiration, GI – gastrointestinal, NR– non reported, RCTs – randomized controlled trials, SNRIs – serotonin–norepinephrine reuptake inhibitors, SARIs – serotonin antagonist and reuptake inhibitors, SSRIs – selective serotonin reuptake inhibitors, TCAs – tricyclic antidepressants, MAOIs – monoamine oxidase inhibitors, NSAIDs – Nonsteroidal anti-inflammatory drugs, UGIB – upper gastrointestinal bleeding.

eTable 3 Suggestive evidence (Class III) for the association of antidepressant use and risk of adverse health outcomes in meta-analyses of observational studies

Adverse health outcomes (author, year)	Exposed/Unexposed	n	Random-effects measure, ES (95% CI)	Results	Criteria used for classification of level of evidence						
					N cases/ Cohort	p- random effects	I ² (p)	PI (95% CI)	SSE/ESB	LS	CE
Pulmonary hypertension (pregnancy maternal exposure; Masarwa, 2019)	SSRI or SNRI users/ No SSRI or SNRI users	11	OR, 1.82 (1.31, 2.53)	Increased risk for SSRIs or SNRI	13 304/ 7 080 850	3.6x10 ⁻⁴	72 (<0.000)	0.73-4.56	No/NP	Yes	III
Pulmonary hypertension (late pregnancy exposure; Masarwa, 2019)	SSRI or SNRI users/ No SSRI or SNRI users	8	OR, 2.09 (1.44, 3.02)	Increased risk for SSRIs or SNRI	12 678/ 5 979 785	9.2x10 ⁻⁵	76 (<0.000)	0.77-5.68	No/No	Yes	III
Autism spectrum disorders (Halvorsen, 2019)	SSRI users/ No SSRI users	9	OR, 1.40 (1.22, 1.60)	Increased risk for SSRIs	4 334/ 1 2682 40	9.7x10 ⁻⁷	15 (0.307)	1.09-1.79	No/NP	No	III
Attention-deficit hyperactivity disorder (Halvorsen, 2019)	SSRI users/ No SSRI users	5	OR, 1.39 (1.17, 1.66)	Increased risk for SSRIs	21 688/ 1 179 596	2.3x10 ⁻⁴	56 (0.061)	0.82-2.37	No/NP	Yes	III
Attention-deficit hyperactivity disorder in children (Jiang, 2018)	Prenatal exposure to ADs/ No AD users	6	RR, 1.34 (1.14,1.57)	Increased risk for ADs	56 272/ 2 840 554	3.3x10 ⁻⁴	79 (<0.000)	0.81–2.21	No/NP	Yes	III
Attention-deficit hyperactivity disorder in children (2+3 trimester; Jiang, 2018)	Prenatal exposure to ADs/ No AD users	5	RR, 1.37 (1.17,1.60)	Increased risk for ADs	41 564/ 2 281 198	6.9x10 ⁻⁵	0 (0.754)	1.06-1.76	No/NP	Yes	III
Attention-deficit hyperactivity disorder in children (Man, 2018)	Pre-conception exposure to ADs/ Non-AD users	5	RR, 1.56 (1.24, 1.96)	Increased risk for ADs	40147/ 1864720	1.4x10 ⁻⁴	58 (0.051)	0.75-3.22	No/NP	Yes	III
Cataract development (Fu, 2018)	SSRIs/non-users or no users of any other AD	6	OR, 1.12 (1.06, 1.19)	Increased risk for SSRIs	446956/ 1 955 042	8.8x10 ⁻⁵	92 (<0.000)	0.92-1.37	Yes/Yes	Yes	III
Cardiovascular malformations (Zhang, 2017)	Any SSRI users/ Non-SSRI users	19	RR, 1.26 (1.13, 1.39)	Increased risk for SSRIs	75362/ 7 368 339	2.1x10 ⁻⁵	54 (0.003)	0.92-1.72	No/NP	Yes	III
Both atrial and ventricular septal defect (Zhang, 2017)	Any SSRI users/ Non-SSRI users	18	RR, 1.27 (1.14, 1.42)	Increased risk for SSRIs	45247/ 10139043	1.6x10 ⁻⁵	40 (0.041)	0.95-1.71	No/NP	Yes	III

Septal defects (Gao, 2017)	Fluoxetine/ No ADs or teratogens use	8	RR, 1.38 (1.19, 1.61)	Increased risk for fluoxetine	39987/ 6438941	2.7×10^{-5}	0 (0.891)	1.14-1.67	No/NP	Yes	III
Cerebrovascular disease (Biffi, 2017)	SSRIs/ Non-SSRIs users	6	RR, 1.26 (1.14, 1.39)	Increased risk for SSRIs	3204/ 280784	1.0×10^{-5}	14 (0.332)	1.02-1.55	No/No	Yes	III
Preterm birth (unadjusted estimates; Eke, 2016)	SSRIs/ Non-SSRIs users	8	OR, 1.59 (1.31, 1.92)	Increased risk for SSRIs	69 912/ 1 307 761	2.0×10^{-5}	92 (<0.000)	0.91-2.78	Yes/Yes	Yes	III
Respiratory distress syndrome (Eke, 2016)	SSRIs/ Non-SSRIs users	5	OR, 1.33 (1.14, 1.55)	Increased risk for SSRIs	19032/ 1269710	2.6×10^{-4}	83 (<0.000)	0.83-2.12	Yes/Yes	Yes	III
Major malformations (Bérard, 2016)	Paroxetine/ No paroxetine use	15	OR, 1.23 (1.10, 1.38)	Increased risk for paroxetine	26752/ 2 169 318	3.8×10^{-4}	3 (0.424)	1.06-1.43	No/NP	No	III
Cardiac malformations (Bérard, 2016)	Paroxetine/ No paroxetine use	18	OR, 1.28 (1.11, 1.47)	Increased risk for paroxetine	75953/ 5109058	2.3×10^{-5}	0 (0.653)	1.09-1.49	No/NP	No	III
Requirement of blood/RBC transfusion (Singh, 2015)	SSRIs+SNRIs/ Non SADs users	7	OR, 1.19 (1.09, 1.30)	Increased risk for SADs	79775/ 556120	1.0×10^{-4}	36 (0.155)	0.97-1.46	No/No	Yes	III
Upper gastrointestinal bleeding (Jiang, 2015)	SSRIs/ Non-SSRIs users	8	1.95 (1.44, 1.93)	Increased risk for SSRIs	32830/ 241266	1.2×10^{-5}	90 (<0.000)	0.70-5.38	No/NP	Yes	III
Low birth weight (Huang, 2014)	Any AD users/ Non- AD users	15	RR, 1.44 (1.21, 1.70)	Increased risk for ADs	20190/ 3001141	2.8×10^{-5}	61 (0.001)	0.88-2.32	No/NP	Yes	III
Preterm birth during late pregnancy (Huybrechts, 2015)	Any AD users/ Non- AD users	12	OR, 1.98 (1.65, 2.37)	Increased risk for ADs	NR/ 1891969	2.4×10^{-13}	83 (<0.000)	1.09-2.37	Yes/NP	Yes	III
Gestational age (Ross, 2013)	Any AD users/ Non- AD users	16	SMD, -0.23 (-0.33, -0.12)	Increased risk for ADs	3419/ 60644	3.9×10^{-5}	71 (<0.000)	-0.81, 0.36	No/Yes	Yes	III

RBC–red blood cells, AD– antidepressants, TCAs – tricyclic antidepressants, SSRIs – selective serotonin reuptake inhibitors, SNRIs – serotonin–norepinephrine reuptake inhibitors, n – number of included studies per association, ES – effect size, N – number of cases, I^2 –heterogeneity, PI – prediction interval, CI – confidence interval, SSE – small-study effect, ESB –excess significance bias, LS – largest study with significant effect, CE – class of evidence, OR – odds ratio, RR – relative risk, SMD – standardized mean difference, NA– not applicable, NP – not pertinent because the number of observed studies is less than the expected.

eTable 4 Weak evidence (Class IV) for the association of antidepressant use and risk of adverse health outcomes in meta-analyses of observational studies

Adverse health outcomes (author, year)	Exposed/Unexposed	n	Random-effects measure, ES (95% CI)	Results	Criteria used for classification of level of evidence						
					N cases/ Cohort	p- random effects	I ² (p)	PI (95% CI)	SSE/ESB	LS	CE
Mental retardation (Halvorsen, 2019)	SSRI users/ No SSRI users	2	HR, 1.41 (1.03, 1.91)	Increased risk for SSRIs	NR/ NR	0.030	0 (0.718)	NA	NA/NA	No	IV
Dementia (Wang, 2018)	SSRI users/ No SSRI users	5	RR, 1.75 (1.03, 2.96)	Increased risk for SSRIs	NR / 53955	0.037	98 (<0.000)	0.23–13.49	No/NA	Yes	IV
Dementia (Wang, 2018)	TCA users/ No SSRI users	4	RR, 2.13 (1.43, 3.18)	Increased risk for TCAs	NR/ 22768	2.1x10 ⁻⁴	96 (<0.000)	0.34-13.54	No/NA	Yes	IV
Dementia (Wang, 2018)	MAOI users/ No MAOI users	2	RR, 2.79 (1.09, 7.17)	Increased risk for MAOIs	NR/ 12 209	0.033	80 (0.025)	NA	NA/NA	Yes	IV
Attention-deficit hyperactivity disorder (first trimester; Jiang, 2018)	Prenatal exposure to ADs/ No AD users	4	RR, 1.28 (1.00,1.64)	Increased risk for ADs	55988/ 293431	0.047	91 (<0.000)	0.41–4.01	No/NP	Yes	IV
Attention-deficit hyperactivity disorder in children (Jiang, 2018)	Maternal psychiatric disorder without exposure versus no exposure	3	RR, 1.34 (1.08,1.67)	Increased risk for ADs	NR/ NR	0.007	60 (0.083)	0.13-13.88	No/NA	Yes	IV
Attention-deficit hyperactivity disorder in children (Jiang, 2018)	SSRIs / Non-SSRIs users	5	RR, 1.35 (1.13,1.61)	Increased risk for ADs	NR/ NR	0.001	86 (<0.000)	0.72-2.51	No/NA	Yes	IV
Attention-deficit hyperactivity disorder in children (Jiang, 2018)	Non-SSRI antidepressant use/ Non-AD users	6	RR, 1.49 (1.22,1.83)	Increased risk for ADs	NR/ NR	1.2x10 ⁻⁴	0 (0.790)	1.12-1.99	No/NA	Yes	IV
Gestational hypertension or preeclampsia (Guan, 2018)	SSRI users/ No SSRI users	7	RR, 1.21 (1.05, 1.40)	Increased risk for SSRIs	12097/ 1108261	0.007	71 (0.002)	0.81-1.82	Yes/Yes	No	IV
Atrial septal defects (Gao, 2018)	Any SSRI users/ Non-SSRI users	7	RR, 1.83 (1.22, 2.72)	Increased risk for SSRIs	6366/ 2560254	0.003	72 (0.002)	0.53-6.31	No/Yes	No	IV
Septal defects (Gao, 2018)	Any SSRI users/ Non-SSRI users	6	RR, 1.38 (1.00, 1.91)	Increased risk for SSRIs	8029/ 2039943	0.050	67 (0.009)	0.50-3.80	No/NP	Yes	IV
Right ventricular outflow tract defects	Any SSRI users/ Non-SSRI users	4	RR, 1.38 (1.09, 1.75)	Increased risk for SSRIs	4307/ 2360018	0.007	33 (0.214)	0.63-3.03	No/NP	Yes	IV

(Gao, 2018)											
Gestational hypertension (Guan, 2018)	SSRI users/ No SSRI users	4	RR, 1.14 (1.00-1.30)	Increased risk for SSRIs	4370/ 91 282	0.048	6 (0.365)	0.79-1.64	No/NP	No	IV
Dental implant failure (Chappuis, 2018)	SSRI users/ No SSRI users	2	OR, 2.02 (1.42, 2.88)	Increased risk for SSRIs	341/ 5 332	9.8x10 ⁻⁵	0 (0.770)	NA	NA/No	Yes	IV
GI bleeding (Na, 2018)	Mirtazapine/ no medication	4	OR, 1.17 (1.01,1.37)	Increased risk for mirtazapine	2116/ 59571	0.043	0 (0.593)	0.83-1.65	No/NP	No	IV
Cataract development (Fu, 2018)	SNRIs/ non-users or no users of any other AD	4	OR, 1.13 (1.04, 1.24)	Increased risk for SNRIs	239390/ 646822	0.006	68 (0.026)	0.79-1.63	Yes/No	Yes	IV
Autism spectrum disorders (Zhou, 2018)	Any AD users/ Non- AD users	13	OR, 1.28 (1.07, 1.53)	Increased risk for SSRIs	26695/ 2699647	0.007	62 (0.002)	0.72-2.26	No/NP	No	IV
Cardiovascular malformations (Shen, 2017)	Sertraline users/Non- sertraline users	12	OR, 1.36 (1.06, 1.74)	Increased risk for sertraline	57493/ 6468241	0.015	64 (0.001)	0.66-2.79	No/NP	Yes	IV
Non-septal defects (Gao, 2017)	Fluoxetine/ No ADs or teratogens use	9	RR, 1.39 (1.12, 1.73)	Increased risk for fluoxetine	14240/ 10822445	0.003	7 (0.374)	0.98-1.97	No/NP	Yes	IV
Preterm birth (adjusted) (Eke, 2016)	SSRIs/ Non-SSRIs users	3	OR, 1.24 (1.09, 1.41)	Increased risk for SSRIs	66652/ 967061	0.001	73 (0.026)	0.31-4.96	No/NP	Yes	IV
Major malformations (Bérard, 2016)	Paroxetine/ No any other SSRIs use	14	OR, 1.19 (1.06, 1.34)	Increased risk for paroxetine	82965/ 4052678	0.003	0 (0.712)	1.05-1.35	No/NP	No	IV
Major malformations (Bérard, 2016)	Paroxetine/ No any other ADs use	10	OR, 1.19 (1.05, 1.35)	Increased risk for paroxetine	100735/ 5766472	0.008	0 (0.578)	1.02-1.38	No/NP	No	IV
Cardiac malformations (Bérard, 2016)	Paroxetine/No any other SSRIs use	17	OR, 1.27 (1.10,1.47)	Increased risk for paroxetine	75138/ 5103306	0.001	0 (0.596)	1.08-1.49	No/NP	No	IV
Cardiac malformations (Bérard, 2016)	Paroxetine/ No any other ADs use	13	OR, 1.23 (1.06, 1.43)	Increased risk for paroxetine	102941/ 6435313	0.006	1 (0.436)	1.03-1.48	No/NP	No	IV
Autistic Spectrum or related Disorders (Healy, 2016)	SSRIs / Non-SSRIs users	17	OR, 1.96 (1.33, 2.88)	Increased risk for SSRIs	23641/ 2321521	0.001	91 (<0.000)	0.42-9.02	No/NP	Yes	IV
Ischemic stroke (Shin, 2014)	SSRIs / Non-SSRIs users	6	OR, 1.48 (1.08, 2.02)	Increased risk for SSRIs	11080/ 724936	0.015	84 (<0.000)	0.50-4.35	No/No	No	IV

Hemorrhagic stroke (Shin, 2014)	SSRIs / Non-SSRIs users	11	OR, 1.32 (1.02, 1.71)	Increased risk for SSRIs	11513/399305	0.033	75 (<0.000)	0.57-3.07	No/NP	Yes	IV
All types of stroke (Shin, 2014)	SSRIs / Non-SSRIs users	6	OR, 1.40 (1.09, 1.80)	Increased risk for SSRIs	3519/223986	0.008	93 (<0.000)	0.57-3.44	No/Yes	Yes	IV
Coronary heart disease (Oh, 2014)	TCAs/ Non-TCAs users	14	OR, 1.51 (1.07, 2.12)	Increased risk for TCAs	6443/347750	0.019	97 (<0.000)	0.41-5.56	No/NP	Yes	IV
Apgar score at 1 minute (Ross, 2013)	Any AD users/ Non-AD users	11	SMD,-0.19 (-0.30,-0.08)	Increased risk for ADs	714/1534	0.001	7 (0.376)	-0.72, 0.34	No/NP	No	IV
Cardiovascular malformations (Riggin, 2013)	Fluoxetine/ Non fluoxetine or teratogens users	16	OR, 1.60 (1.32, 1.95)	Increased risk for fluoxetine	35702/3401555	2.4×10^{-6}	1 (0.441)	1.27-2.02	No/NP	No	IV
Poor neonatal adaptation (Grigoriadis, 2013)	Any AD users/ Non-AD users	8	OR, 5.06 (3.25, 7.89)	Increased risk for ADs	75/986	7.8×10^{-13}	0 (0.617)	2.91-8.81	No/No	Yes	IV
Respiratory distress syndrome (Grigoriadis, 2013)	Any AD users/ No AD users	9	OR, 2.20 (1.81, 2.66)	Increased risk for ADs	623/583525	1.3×10^{-15}	38 (0.115)	1.43-3.37	Yes/Yes	Yes	IV
Tremors (Grigoriadis, 2013)	Any AD users/ Non-AD users	4	OR, 7.90 (3.33, 18.73)	Increased risk for ADs	60/482	2.7×10^{-6}	45 (0.144)	0.35-177.44	No/Yes	Yes	IV
Major malformations (Nikfar, 2013)	SSRIs / Non-SSRIs users	21	OR, 1.23 (1.08, 1.41)	Increased risk for SSRIs	60753/1761861	0.002	25 (0.146)	0.91-1.68	Yes/No	No	IV
Spontaneous abortion (Nikfar, 2013)	SSRIs / Non-SSRIs users	7	OR, 1.85 (1.41, 2.42)	Increased risk for SSRIs	376/4140	9.6×10^{-6}	3 (0.403)	1.25-2.74	No/No	No	IV
Breast and ovarian cancer (Cosgrove, 2011)	SSRIs / Non-SSRIs users	16	OR, 1.05 (1.01, 1.10)	Increased risk for SSRIs	46796/1059784	0.010	15 (0.287)	0.98-1.14	Yes/No	No	IV
Suicide attempt and completion in adults (Barbui, 2009)	SSRIs / Non-SSRIs users	2	OR, 0.47 (0.27, 0.80)	Decreased risk for SSRIs	107/178529	4.1×10^{-9}	0 (0.555)	NA	NA/NP	No	IV
Spontaneous abortion (Hemels, 2005)	Any AD users/ Non-AD users	11	OR, 1.46 (1.19, 1.79)	Increased risk for ADs	373/3194	2.8×10^{-3}	0 (0.981)	1.15-1.85	No/NP	No	IV

AD–antidepressants, APGAR – backronym of Appearance (skin color), Pulse (heart rate), Grimace (reflex irritability), Activity (muscle tone), and Respiration, TCAs – tricyclic antidepressants, SSRIs – selective serotonin reuptake inhibitors, SNRIs – serotonin–norepinephrine reuptake inhibitors, n – number of included studies per association, ES – effect size, N – number of cases, I^2 –heterogeneity, PI – prediction interval, CI – confidence interval, SSE – small-study effect, ESB –excess significance bias, LS – largest study with

significant effect, CE – class of evidence, OR – odds ratio, RR – relative risk, NA– not applicable, NP – not pertinent because the number of observed studies is less than the expected.

eTable 5 No evidence (Non-significant associations, $p>0.05$) for the association of antidepressant use and risk of adverse health outcomes in meta-analyses of observational studies

Adverse health outcomes (author, year)	Exposed/ Unexposed	n	Random-effect measure, ES (95% CI)	Results	Criteria used for classification of level of evidence						
					N cases/ Cohort	p- random effects	I ² (p)	PI (95% CI)	SSE/ES B	LS	CE
Major congenital anomalies (Gao, 2018)	Maternal psychiatric disorder with SSRI exposure versus no exposure	4	RR, 1.03 (0.95, 1.13)	No evidence of risk	63383/ 1851983	0.439	3 (0.380)	0.85-2.26	No/No	No	NS
Cardiovascular malformations (Gao, 2018)	Maternal psychiatric disorder with SSRI exposure versus no exposure	6	RR, 1.06 (0.89, 1.26)	No evidence of risk	22192/ 2685027	0.489	34 (0.182)	0.71-1.60	No/NP	No	NS
Left ventricular outflow tract defects (Gao, 2018)	SSRIs / Non-SSRIs users	3	RR, 1.08 (0.81, 1.44)	No evidence of risk	3117/ 1410514	0.616	0 (0.892)	0.17-6.99	No/NP	No	NS
Ventricular septal defect (Gao, 2018)	SSRIs / Non-SSRIs users	8	RR, 1.10 (0.93, 1.29)	No evidence of risk	13647/ 3509759	0.232	36 (0.140)	0.76-1.60	NA/NP	Yes	NS
BMD Lumbar Spine (Schweiger, 2018)	SSRIs / Non-SSRIs users	2	SMD, -0.03 (-0.43, 0.38)	No evidence of risk	NR/ 2100	0.897	71 (0.065)	NA	NA/No	Yes	NS
BMD Femoral Neck (Schweiger, 2018)	TCAs / Non-TCAs users	2	SMD, 0.02 (-0.14, 0.18)	No evidence of risk	NR/ 3851	0.786	0 (0.545)	NA	NA/NP	No	NS
BMD Total Hip (Schweiger, 2018)	SSRIs / Non-SSRIs users	2	SMD, 0.10 (-0.38, 0.59)	No evidence of risk	NR/ 4694	0.677	98 (<0.000)	NA	NA/No	Yes	NS
BMD Total Hip (Schweiger, 2018)	TCAs / Non-TCAs users	2	SMD, 0.03 (-0.12, 0.17)	No evidence of risk	NR/ 4694	0.719	0 (0.644)	NA	NA/NP	No	NS
Attention-deficit hyperactivity disorder in children (Jiang, 2018)	Exposure versus maternal psychiatric disorder without exposure	2	RR, 0.96 (0.76,1.20)	No evidence of risk	NR/ NR	0.714	0 (0.524)	NA	NA/NA	No	NS
Attention-deficit hyperactivity disorder in children (Jiang, 2018)	Sibling matched	3	RR, 0.88 (0.70,1.11)	No evidence of risk	NR/ NR	0.293	0 (0.486)	0.20-3.99	Yes/NA	No	NS

Preeclampsia (Guan, 2018)	SSRI users/ No SSRI users	5	RR, 1.32 (0.98-1.78)	No evidence of risk	7 727/ 201 181	0.071	83 (<0.000)	0.47-3.70	No/No	No	NS
GI bleeding (Na, 2018)	Mirtazapine/SSRIs	4	OR, 1.03 (0.89,1.19)	No evidence of risk	2116/ 59571	0.689	0 (0.806)	0.75 -1.41	No/NP	No	NS
Any other type of bleeding (Na, 2018)	Bupropion/SSRIs	3	OR, 0.90 (0.68,1.18)	No evidence of risk	1228/ 16223	0.443	0 (0.886)	0.15-5.24	No/NP	No	NS
Both GI and any other type of bleeding (Na, 2018)	Mirtazapine/SSRIs	5	OR, 0.98 (0.98,1.14)	No evidence of risk	2751/ 63545	0.766	6 (0.371)	0.77-1.25	No/NP	No	NS
Both GI and any other type of bleeding (Na, 2018)	Mirtazapine/ no medication	4	OR, 1.12 (0.97,1.29)	No evidence of risk	2116/ 59571	0.131	0 (0.619)	0.81-1.53	No/NP	No	NS
Transfusion of FFP (Eckersley, 2018)	SSRIs users/ Non- SSRI users	3	OR, 0.96 (0.74,1.24)	No evidence of risk	8510/ 139711	0.754	60 (0.083)	0.06-14.93	No/NP	No	NS
Transfusion of platelets (Eckersley, 2018)	SSRIs users/ Non- SSRI users	3	OR, 0.93 (0.78, 1.11)	No evidence of risk	4424/ 139711	0.418	5 (0.349)	0.26-3.28	No/NP	No	NS
Re-operation for bleeding (Eckersley, 2018)	SSRIs users/ Non- SSRI users	3	OR, 1.07 (0.66, 1.74)	No evidence of risk	446/ 11955	0.783	0 (0.964)	0.05-25.20	No/NP	No	NS
Thirty-day mortality (Eckersley, 2018)	SSRIs users/ Non- SSRI users	5	OR, 1.03 (0.91, 1.16)	No evidence of risk	6668/ 270103	0.664	0 (0.814)	0.84-1.26	No/NP	No	NS
Autism spectrum disorders (Morales, 2018)	Any AD users/ Non-AD users with a history of affective disorder	6	RR, 1.18 (0.91, 1.52)	No evidence of risk	7780/ 1052964	0.192	51 (0.069)	0.58 -2.40	No/NP	No	NS
Nervous system (Shen, 2017)	Sertaline users/Non- sertraline users	6	OR, 1.43 (0.88, 2.32)	No evidence of risk	416/ 1975394	0.149	0 (0.976)	0.72-2.84	No/NP	No	NS

Digestive system (Shen, 2017)	Sertaline users/Non-sertraline users	5	OR, 1.23 (0.76, 1.98)	No evidence of risk	720/1657603	0.407	0 (0.805)	0.56-2.68	No/NP	No	NS
Eye, ear, face and neck (Shen, 2017)	Sertaline users/Non-sertraline users	4	OR, 1.08 (0.33, 3.55)	No evidence of risk	316/1839470	0.902	32 (0.219)	0.02-56.42	No/NP	No	NS
Urogenital system (Shen, 2017)	Sertaline users/Non-sertraline users	8	OR, 1.03 (0.73,1.46)	No evidence of risk	1638/1676096	0.869	0 (0.755)	0.67-1.59	Yes/NP	No	NS
Musculoskeletal system (Shen, 2017)	Sertaline users/Non-sertraline users	5	OR, 0.97 (0.69, 1.36)	No evidence of risk	1694/1657603	0.861	0 (0.723)	0.56-1.68	No/NP	No	NS
Nervous system malformations (Gao, 2017)	Fluoxetine/ No ADs or teratogens use	3	RR, 1.37 (0.83, 2.25)	No evidence of risk	2925/1817322	0.219	0 (0.526)	0.05-34.60	No/NP	No	NS
Eye malformations (Gao, 2017)	Fluoxetine/ No ADs or teratogens use	3	RR, 1.30 (0.53, 3.17)	No evidence of risk	2585/1270645	0.564	0 (0.399)	0.00-420.44	No/NP	No	NS
Urogenital malformations (Gao, 2017)	Fluoxetine/ No ADs or teratogens use	5	RR, 1.02 (0.65, 1.59)	No evidence of risk	4244/2262620	0.932	39 (0.159)	0.00-3.50	No/NP	Yes	NS
Digestive malformations (Gao, 2017)	Fluoxetine/ No ADs or teratogens use	3	RR, 1.08 (0.60, 1.96)	No evidence of risk	2019/1816416	0.800	0 (0.861)	0.02-51.29	No/NP	No	NS
Respiratory malformations (Gao, 2017)	Fluoxetine/ No ADs or teratogens use	3	RR, 1.38 (0.69, 2.78)	No evidence of risk	472/1814869	0.361	0 (0.670)	0.02-127.68	No/NP	No	NS
Musculoskeletal malformations (Gao, 2017)	Fluoxetine/ No ADs or teratogens use	4	RR, 0.82 (0.54, 1.22)	No evidence of risk	1474/1910723	0.322	0 (0.726)	0.34-1.98	Yes/NP	No	NS

Acute heart disease (Biffi, 2017)	Any AD users/ Non-AD users	7	RR, 1.35 (0.91, 2.02)	No evidence of risk	65331/ 818933	0.138	92 (<0.000)	0.35-5.19	No/NP	Yes	NS
Acute heart disease (Biffi, 2017)	SSRIs/ Non-SSRIs users	14	RR, 1.00 (0.83,1.22)	No evidence of risk	89421/ 818337	0.961	85 (<0.000)	0.53-1.90	Yes/NP	Yes	NS
Cerebrovascular disease (Biffi, 2017)	TCAs/ Non-TCAs users	4	RR, 1.06 (0.96, 1.17)	No evidence of risk	7325/ 278749	0.239	0 (0.745)	0.85-1.32	No/NP	No	NS
Preterm birth (Huybrechts, 2015)	Any AD users in early pregnancy/ Non-AD users	8	OR, 1.15 (0.96, 1.38)	No evidence of risk	NR/ 1371456	0.127	85 (<0.000)	0.64-2.07	No/NP	Yes	NS
Myocardial infarction (Undela, 2015)	Any AD users/ Non-AD users	21	RR, 1.03 (0.88, 1.22)	No evidence of risk	220362/ 1793877	0.687	98 (<0.000)	0.51-2.08	No/NP	Yes	NS
Requirement of reoperation for bleeding complication (Singh, 2015)	SSRIs+SNRIs/ Non SADs users	4	OR, 1.48 (0.84, 2.62)	No evidence of risk	676/ 26743	0.176	54 (0.089)	0.17-13.06	No/NP	Yes	NS
Mortality-any cause (Singh, 2015)	SSRIs+SNRIs/ Non SADs users	5	OR, 1.15 (0.86, 1.53)	No evidence of risk	79683/ 554079	0.343	76 (0.002)	0.46-2.88	No/No	Yes	NS
Requirement of RBC transfusion (Singh, 2015)	Any other AD users/ Non-any other AD users	3	OR, 1.03 (0.74, 1.43)	No evidence of risk	216/ 5495	0.874	0 (0.790)	0.12-8.88	No/NP	No	NS
Pulmonary hypertension of the newborn (Grigoriadis, 2014)	SSRIs early in pregnancy/ Non- SSRIs users	3	OR,1.22 (0.58, 2.60)	No evidence of risk	49/ 13017	0.600	78 (0.010)	0.00-7225.40	No/NP	No	NS
Major malformations (Riggin, 2013)	Fluoxetine/ No fluoxetine or teratogens use	22	OR,1.12 (0.98, 1.28)	No evidence of risk	163333/ 4576977	0.109	29 (0.106)	0.78-1.61	No/NP	No	NS
Cardiovascular malformations (Myles, 2013)	Citalopram during first trimester / Non-AD users	6	OR, 0.99 (0.75, 1.30)	No evidence of risk	NR/NR	0.459	34 (0.183)	0.51-1.92	No/NA	No	NS

Major malformations (Myles, 2013)	Citalopram during first trimester Non-AD users	7	OR, 1.04 (0.92, 1.17)	No evidence of risk	NR/NR	0.254	0 (0.858)	0.89-1.21	No/NA	No	NS
Breast cancer (Eom, 2012)	Any AD users/ Non-AD users	18	OR, 1.02 (0.96, 1.08)	No evidence of risk	65313/ 739891	0.574	36 (0.063)	0.87-1.18	Yes/NP	No	NS
Colon cancer (Lee, 2012)	SSRIs / Non-SSRIs users or any other ADs	6	OR, 0.89 (0.79, 1.01)	No evidence of risk	11710/ 978578	0.071	41 (0.131)	0.66-1.21	No/No	No	NS
Breast and ovarian cancer (Cosgrove, 2011)	TCAs/Non-TCAs	16	OR, 1.03 (0.98, 1.09)	No evidence of risk	37923/ 426545	0.251	47 (0.021)	0.88-1.22	No/NP	No	NS

GI – gastrointestinal, RBC–red blood cells, FFP– fresh frozen plasma, BMD – bone mineral density, AD– antidepressants, TCAs – tricyclic antidepressants, SSRIs – selective serotonin reuptake inhibitors, SNRIs – serotonin–norepinephrine reuptake inhibitors, SADs– serotonergic antidepressants, n – number of included studies per association, ES – effect size, N – number of cases, I² –heterogeneity, PI – prediction interval, CI – confidence interval, SSE – small-study effect, ESB –excess significance bias, LS – largest study with significant effect, CE – class of evidence, CES= OR – odds ratio, RR – relative risk, HR– hazard ratio, SMD – standardized mean difference, NA– not applicable, NP – not pertinent because the number of observed studies is less than the expected.

Supplementary methods

Selection between overlapping meta-analysis

At full-text assessment, we extracted relevant information to define antidepressants exposure, inclusion criteria for the population of interest and outcome definition of each meta-analysis. When two or more meta-analyses focused on the same combination of exposure, population, and outcome we selected only the meta-analysis that included the largest data set and sufficient individual data for statistical analysis.²⁰⁸ We also examined whether the main reported conclusions were concordant regarding the direction and the significance level of the association.² If the results were concordant, then again, we selected the one that included the largest data set and sufficient individual data for statistical analysis.²⁰⁸ If two or more meta-analyses focused on the same combination of exposure, population, and outcome, but included a different set of primary studies (for example, different set of cohort studies but the same set of case-control studies and vice versa, or only cohort studies) then we kept both or all. If two or more meta-analyses focused on the same combination of exposure, population, and outcome and included exactly the same number and set of included studies then we kept the most recent one.²⁰⁹ We adopted this procedure not only to avoid overlapping data sets as much as possible, but also not to miss as much any valuable information. We also examined the concordance between selected and non-selected meta-analyses by sensitivity analysis limited to non-selected associations due to overlap (eTable 6).²¹⁰

Quality assessment

AMSTAR 2²¹¹ assesses whether included meta-analyses clearly defined the research question and inclusion criteria, mentioned an a-priori protocol, explained inclusion criteria, conducted a comprehensive literature search, screened literature and extracted data in duplicate, provided a list of excluded studies with reason for exclusion, described included studies in detail, assessed risk of bias of included studies, reported on the source of funding, used appropriate statistical methods, accounted for risk of bias for instance with sensitivity analyses, considered risk of bias when interpreting results, explained and discussed any heterogeneity in results, assessed and discussed

publication bias, reported any potential conflict of interest. Based on the above listed items, AMSTAR 2 defines a systematic review's quality as high, moderate, low, or critically low. Compared with AMSTAR, AMSTAR 2 also assesses the quality of the studies included in a meta-analysis, without limiting the quality assessment to the technical aspects of the meta-analysis itself. Additionally, compared to ROBIS, AMSTAR has much better agreement among raters.²¹²

Grading method

We assessed the credibility of the evidence using established criteria published in several umbrella reviews.^{208,209,210} Specifically, associations had the strongest validity and were not suggestive of bias (Class I: convincing) whenever they met all the following criteria: had >1000 cases; had p-value <10⁻⁶ based on random-effects meta-analysis; had no evidence of small-study effects and excess significance; had 95% prediction interval (PI) that excluded the null value; the largest study had a nominally significant effect (p<0.05); and had low or moderate between-study heterogeneity (I² <50%). Highly suggestive evidence (Class II) required >1000 cases; had p-value <10⁻⁶ based on random-effects meta-analysis; and the largest study had a nominally significant effect (p<0.05). Suggestive evidence (Class III) criteria required only >1000 cases and had p-value <10⁻³ based on random-effects meta-analysis. Weak evidence (Class IV) criteria required only all other remained associations with a p-value ≤0.05. These criteria have been proven to provide a systematic and transparent judgment of the methodological flaws that can occur in various meta-analyses. It is well known, that heterogeneity, publication bias, small-study effects, selective reporting, and excess of significant bias in the published meta-analyses can contribute to biased results of a meta-analyses. Thus, this method appears to be more objective than other quality grading methods because it uses several statistical tests to assess different type of bias and it can work for many types of research questions.²¹³ Notably, an empirical evaluation of different grading approaches concluded that agreement was poor and that all methods had important shortcomings.²¹⁴

Supplementary results

Sensitivity analysis results of convincing and highly suggestive evidence associations from the main analysis

Cohort studies (both prospective and retrospective)

A sensitivity analysis limited to cohort studies showed that none of the associations within class I retained the same rank (Table 3). However, this analysis showed that six associations were supported by highly suggestive evidence (Table 3). These included the increased risk of autism during pregnancy, osteoporotic fractures, pre-term birth and lower APGAR scores at 5 minutes (antidepressants during pregnancy), and suicide attempt/completion in children and adolescents, as well as the decreased risk of suicide attempt/completion in adults and older adults.

The associations between antidepressants and attention-deficit hyperactivity disorder in children, postpartum haemorrhage, and osteoporotic fractures (TCAs) were downgraded to suggestive evidence, while the associations between antidepressants and ASD during pre-pregnancy, severe bleeding, upper gastrointestinal bleeding, and hip fracture were downgraded to weak evidence.

Prospective cohort studies

A sensitivity analysis limited to prospective cohort studies showed that none of the associations within class I retained the same rank (Table 3). The most important change was for highly suggestive associations, with one being upgraded to convincing evidence (preterm birth related to any antidepressant use), while all other were downgraded to lower ranks of evidence. More analytically, the associations between antidepressants and attention-deficit hyperactivity disorder in children, and postpartum haemorrhage were downgraded to suggestive evidence, while the associations with severe bleeding and Apgar score at 5 minutes were downgraded to weak evidence. Finally, the association with autism during pre-pregnancy turned into non-significant.

Studies adjusted for multiple covariates

When the sensitivity analysis was limited to studies adjusted for multiple potential confounders beyond age and sex (as described in eTable 7 in the Supplement), the association with suicide risk

in children and adolescents, the association between any antidepressant and ASD during pre-pregnancy remained at convincing evidence, while five more associations remained at highly suggestive evidence as in the main analysis (Table 5). These included, increased risk of attention-deficit hyperactivity disorder in children, osteoporotic fractures (SSRIs and TCAs), hip fracture (any antidepressant), and upper gastrointestinal bleeding. The positive associations with cataract development, severe bleeding, and postpartum haemorrhage were downgraded to suggestive evidence as well as the protective association with suicide risk in adults, whereas the association with preterm birth, and Apgar score at 5 minutes were downgraded to weak evidence. Finally, the association with autism during pregnancy included only unadjusted estimates and excluded from this analysis.

Confounding by indication

This analysis included only adjusted studies that were controlled/matched for a psychiatric condition as depicted in eTable 7 in the Supplement. This analysis showed that none of the associations within class I retained the same rank (Table 3). The association with autism during pre-pregnancy was downgraded to highly suggestive, while the association with suicide attempt/completion in children and adolescents was downgraded to weak (Table 4 in the main text). Two more associations, i.e., osteoporotic fractures (SSRIs and TCAs) remained at highly suggestive evidence as in main analysis (Table 3). The associations with attention-deficit hyperactivity disorder in children, postpartum haemorrhage, and the protective association with suicide risk in adults, were downgraded to suggestive evidence, while the association with preterm birth was downgraded to weak evidence.

High-quality primary studies

This analysis limited to high-quality primary studies that had a Newcastle-Ottawa Scale (NOS) score bigger than seven (or defined as such by using other instruments) as reported from the original meta-analyses' authors. This analysis showed that the three associations within class I retained the same rank (Table 3). These included the increased risk of autism before and during

pregnancy, and suicide attempt and completion in children and adolescents. Seven more associations remained at highly suggestive evidence as in main analysis (Table 3). These included the attention-deficit hyperactivity disorder in children, osteoporotic fractures (SSRIs and TCAs), hip fracture (any antidepressant), postpartum haemorrhage, preterm birth, and the protective association with suicide risk in adults. The remaining associations with cataract development, severe bleeding, upper gastrointestinal bleeding, and Apgar score at 5 minutes were downgraded to weak evidence.

Classes of antidepressants

A sensitivity analysis limited to SSRIs showed that the associations with increased suicide risk in children and adolescents and autism during pregnancy remained convincing, while another one was upgraded to convincing evidence (lower APGAR scores at 5 minutes). Three more associations were supported by highly suggestive evidence (Table 3). These included the osteoporotic fractures, preterm birth, and the protective association with suicide risk in adults.

When the sensitivity analysis was limited to TCAs, two associations remained at highly suggestive evidence i.e., cataract development and osteoporotic fractures.

The analysis limited to other or mixed antidepressants showed that the association between any antidepressant and autism during pre-pregnancy remained convincing, one association was upgraded to convincing evidence (preterm birth), while one was downgraded to suggestive evidence (attention-deficit hyperactivity disorder in children). Another one turned into non-significant, i.e, APGAR scores at 5 minutes. Finally, three other associations remained at highly suggestive evidence (Table3). These included severe bleeding, postpartum haemorrhage, and upper gastrointestinal bleeding.

Studies located in Europe

When the sensitivity analysis was limited to studies located in Europe, the association between SSRIs and autism during pregnancy remained at convincing evidence (Table3), while the association with suicide attempt/completion in children and adolescents was downgraded to weak

evidence. Five associations remained at highly suggestive evidence (upper gastrointestinal bleeding, preterm birth, osteoporotic fractures (SSRIs and TCAs), and hip fractures (either TCAs or SSRIs). The association with severe bleeding and the protective association with suicide risk in adults were downgraded to suggestive evidence, while the associations with attention-deficit hyperactivity disorder in children and Apgar score at 5 minutes were downgraded to weak evidence.

Studies located in North America

When the sensitivity analysis was limited to studies located in North America i.e., USA and Canada, none of the associations within class I retained the same rank (Table 3). Four more associations i.e., osteoporotic fractures (SSRIs and TCAs), hip fractures (either TCAs or SSRIs), and preterm birth remained at highly suggestive evidence as in main analysis (Table 4 in the main text). The associations with autism during pregnancy and severe bleeding were downgraded to suggestive evidence, while the associations with autism before pregnancy, suicide attempt/completion in children and adolescents, attention-deficit hyperactivity disorder in children, postpartum haemorrhage, upper gastrointestinal bleeding, Apgar score at 5 minutes, and the protective association with suicide risk in adults were downgraded to weak evidence.

Studies located in other regions

When the sensitivity analysis was limited to studies located in other regions, none of the associations within class I and II retained the same rank (Table3). The associations with postpartum haemorrhage, upper gastrointestinal bleeding, preterm birth, and hip fracture (either TCAs or SSRIs) were downgraded to weak evidence, while the association with severe bleeding turned into non-significant. For the remaining associations within Class I and II there were no available data.

Sensitivity analysis results of non-selected meta-analyses due to overlap

The results from this analysis are presented in eTable 6. Among the 74 associations, that were overlapped between selected and non-selected meta-analyses with adequate data, only three (4.1%) had convincing evidence; the association between any antidepressant before pregnancy and

SSRI/any antidepressant during pregnancy and autism spectrum disorders (ASD).^{29, 84, 215} Three more associations (4.1%) were supported by highly suggestive evidence. These included the association between SSRI before pregnancy and autism spectrum disorders and osteoporotic fractures.^{57, 137, 151, 13} There was suggestive evidence for eight further associations (10.8%) linked to increased risk of adverse health outcomes. For the rest of the associations, there was either weak (n=32 [43.2%]) or no evidence (n=28 [37.8%]; all associations with $p>0.05$). Overall, we found an agreement between the associations that included in the main analysis and those excluded due to overlap.

eTable 6 Sensitivity analysis results of non-selected meta-analyses due to overlap

Author, year	Risk of adverse health outcome	Antidepressant exposure	Number of cases / total population	Number of study estimates	Study design	Effect metrics	Random effects summary estimate (95% CI)	Random effects summary estimate P value	Largest study summary estimate (95% CI)	I ² (%)	Egger P value	95% prediction interval	Level of evidence
Ng, 2019	Persistent pulmonary hypertension	SSRI during pregnancy	397 / 7470566	9	Cohort, case-control	OR	1.94 (1.37, 2.76)	2.1x10 ⁻³	0.79 (0.08, 7.84)	88	0.95	0.70, 5.39	Weak
Kaplan, 2017	Autism spectrum disorders	Maternal psychiatric disorder without SSRI	100 / 36925	2	Cohort	OR	1.81 (1.44, 2.29)	6.0x10 ⁻⁷	2.02 (1.53, 2.66)	20	0.00	NA	Weak
Mezzacappa, 2017	Autism spectrum disorders	Antidepressant during preconception period	8040 / 69219	4	Case-control	OR	1.77 (1.49, 2.09)	3.3x10 ⁻¹¹	1.90 (1.50, 2.40)	0	0.98	1.22, 2.56	Convincing
Rais, 2014	Autism spectrum disorders	Antidepressant during pregnancy	5040 / 50187	3	Case-control	OR	1.50 (0.74, 3.03)	0.260	2.18 (1.37, 3.46)	84	0.87	0.00, 7910.38	No association
Mezzacappa, 2017	Autism spectrum disorders	Antidepressant during pregnancy	17318 / 889256	7	Cohort, case-control	OR	1.41 (1.12, 1.78)	0.003	1.46 (1.17, 1.82)	63	0.85	0.72, 2.78	Weak
Mezzacappa, 2017	Autism spectrum disorders	Antidepressant during pregnancy	4849 / 772331	2	Cohort	HR	1.26 (0.91, 1.74)	0.160	1.46 (1.17, 1.82)	67	0.00	NA	No association
Mezzacappa, 2017	Autism spectrum disorders	Antidepressant during pregnancy	12469 / 116925	5	Case-control	OR	1.52 (1.09, 2.13)	0.014	1.9 (1.48, 2.44)	61	0.61	0.52, 4.44	Weak
Mezzacappa, 2017	Autism spectrum disorders	Antidepressant during the first trimester	17318 / 889256	7	Cohort, case-control	OR	1.55 (1.19, 2.03)	0.001	1.46 (1.17, 1.82)	68	0.78	0.70, 3.44	Weak
Mezzacappa, 2017	Autism spectrum disorders	Antidepressant during the first trimester	4849 / 772331	2	Cohort	HR	1.26 (0.91, 1.74)	0.161	1.46 (1.17, 1.82)	67	0.00	NA	No association
Mezzacappa, 2017	Autism spectrum disorders	Antidepressant during the first trimester	12469 / 116925	5	Case-control	OR	1.79 (1.27, 2.52)	1.0x10 ⁻³	2.05 (1.58, 2.66)	55	0.76	0.62, 5.13	Suggestive

Mezzacappa, 2017	Autism spectrum disorders	Antidepressant during the second trimester	8040 / 69219	4	Case-control	OR	1.67 (1.14, 2.45)	0.009	2.30 (1.63, 3.24)	38	0.38	0.44, 6.33	Weak
Mezzacappa, 2017	Autism spectrum disorders	Antidepressant during the third trimester	8040 / 69219	4	Case-control	OR	1.54 (0.82, 2.90)	0.180	2.69 (1.81, 4.00)	73	0.55	0.11, 22.57	No association
Kaplan, 2016	Autism spectrum disorders	Non-SSRI antidepressant during pregnancy	3354 / 25728	3	Case-control	OR	2.05 (1.20, 3.49)	0.008	1.93 (0.88, 4.24)	0	0.89	0.07, 64.16	Weak
Kaplan, 2017	Autism spectrum disorders	SSRI discontinuation until 3 months before pregnancy vs. unexposed	3852 / 652201	2	Cohort	RR	1.31 (0.98, 1.74)	0.065	1.46 (1.17, 1.82)	46	0.00	NA	No association
Kaplan, 2016	Autism spectrum disorders	SSRI during preconception period	6890 / 64569	3	Case-control	OR	1.84 (1.48, 2.28)	2.4x10 ⁻⁸	1.90 (1.50, 2.40)	0	0.78	0.46, 7.35	Highly suggestive
Man, 2015	Autism spectrum disorders	SSRI during pregnancy	10424 / 107688	4	Case-control	OR	1.81 (1.47, 2.24)	3.3x10 ⁻⁸	1.80 (1.40, 2.31)	0	0.94	1.14, 2.88	Convincing
Kobayashi, 2016	Autism spectrum disorders	SSRI during pregnancy	10664 / 988245	7	Cohort, case-control	OR	1.45 (1.15, 1.81)	0.002	1.20 (0.90, 1.60)	31	0.18	0.86, 2.43	Weak
Kobayashi, 2016	Autism spectrum disorders	SSRI during pregnancy	4068 / 632851	2	Cohort	OR	1.69 (0.80, 3.56)	0.170	1.20 (0.90, 1.60)	82	0.00	NA	No association
Kobayashi, 2016	Autism spectrum disorders	SSRI during pregnancy	6596 / 355394	5	Case-control	OR	1.38 (1.08, 1.75)	0.009	1.35 (0.94, 1.93)	0	0.38	0.93, 2.03	Weak
Kaplan, 2016	Autism spectrum disorders	SSRI during pregnancy	9061 / 83905	5	Case-control	OR	1.65 (1.23, 2.23)	1.0x10 ⁻³	1.90 (1.47, 2.45)	38	0.76	0.73, 3.73	Suggestive
Brown, 2017	Autism spectrum disorders	SSRI during pregnancy	8695 / 798967	6	Cohort, case-control	OR	1.44 (1.12, 1.85)	0.005	1.20 (0.90, 1.60)	24	0.19	0.82, 2.52	Weak
Brown, 2017	Autism spectrum disorders	SSRI during pregnancy	4849 / 772427	2	Cohort	RR	1.54 (0.86, 2.75)	0.153	1.20 (0.90, 1.60)	70	0.00	NA	No association
Brown, 2017	Autism spectrum disorders	SSRI during pregnancy	3846 / 26540	4	Case-control	OR	1.44 (1.02, 2.02)	0.037	1.00 (0.59, 1.68)	7	0.37	0.62, 3.35	Weak

Kaplan, 2017	Autism spectrum disorders	SSRI during pregnancy	3992 / 679410	3	Cohort	HR	1.61 (1.16, 2.25)	0.005	1.40 (1.02, 1.92)	46	0.33	0.06, 44.64	Weak
Kaplan, 2016	Autism spectrum disorders	SSRI during the first trimester	7382 / 65381	4	Case-control	OR	1.91 (1.28, 2.83)	0.001	2.00 (1.54, 2.60)	42	0.98	0.46, 7.91	Weak
Brown, 2017	Autism spectrum disorders	SSRI during the first trimester	3846 / 26540	4	Case-control	OR	1.75 (1.16, 2.63)	0.007	1.30 (0.72, 2.36)	21	0.64	0.51, 5.95	Weak
Kaplan, 2016	Autism spectrum disorders	SSRI during the second trimester	7382 / 65381	4	Case-control	OR	1.73 (1.15, 2.61)	0.008	2.30 (1.63, 3.25)	29	0.28	0.46, 6.58	Weak
Kaplan, 2016	Autism spectrum disorders	SSRI during the third trimester	7382 / 65381	4	Case-control	OR	1.64 (0.83, 3.24)	0.163	2.70 (1.81, 4.03)	68	0.56	0.10, 26.8	No association
Wang, 2015	Risk of Heart Defects	SSRI during pregnancy	NR/ 2010180	4	Cohort	OR	1.22 (0.89, 1.67)	0.210	1.60 (1.10, 1.30)	92	0.50	0.31, 4.88	No association
McDonagh, 2014	Preterm birth	Antidepressant during pregnancy	NR / 304	2	Cohort, case-control	OR	1.84 (0.79, 4.27)	0.155	1.73 (0.63, 4.58)	0	NA	NA	No association
Anglin, 2014	Upper gastrointestinal bleeding	SSRIs + or NSAIDs (Patients with various disorders with a diagnosis of upper gastrointestinal bleeding)	85628/ 1020230	19	Cohort, case-control	OR	1.66 (1.44, 1.91)	8.2×10^{-13}	1.06 (0.57, 1.97)	82	0.80	0.93, 2.95	Suggestive
Painfully, 2013	Cardiovascular malformations	Paroxetine during first trimester	23746/ 1473816	11	Cohort	RR	1.25 (1.01, 1.54)	0.041	1.03 (0.80, 1.33)	30	0.95	0.78, 2.01	Weak
Grigoriadis, 2013	Major malformations	SSRI during first trimester	NR/ 789337	12	Cohort, case-control	RR	0.93 (0.85, 1.02)	0.121	0.78 (0.45, 1.36)	0	0.26	0.83, 1.03	No association
Grigoriadis, 2013	Cardiovascular malformations	SSRI during first trimester	NR/ 345203	13	Cohort, case-control	RR	1.36 (1.08, 1.71)	0.009	0.50 (0.06, 4.43)	31	0.34	0.81, 2.29	Weak
Wu, 2012	Fracture risk	SSRI exposure in general population of older adults	66050/ 385515	12	Cohort, case-control	RR	1.72 (1.51, 1.95)	4.3×10^{-18}	1.40 (1.34, 1.46)	91	0.06	1.12, 2.61	Highly suggestive
Eom, 2012	Fracture risk	SSRI exposure in general population of older adults	201627/ 916152	12	Cohort, case-control	RR	1.69 (1.51, 1.90)	2.5×10^{-19}	1.40 (1.34, 1.46)	90	0.05	1.14, 2.52	Highly suggestive

Loke, 2008	Upper gastrointestinal bleeding	SSRIs (Patients with various disorders with a diagnosis of upper gastrointestinal bleeding)	2121/127048	4	Cohort, case-control	OR	2.35 (1.44, 3.85)	0.001	3.60 (2.70, 4.70)	96	0.43	0.23, 24.66	Weak
Loke, 2008	Upper gastrointestinal bleeding	SSRIs + NSAIDs (Patients with various disorders with a diagnosis of upper gastrointestinal bleeding)	619/127048	4	Cohort, case-control	OR	6.33 (3.39, 11.81)	6.5x10 ⁻⁹	12.20 (7.10, 19.50)	91	0.16	0.36, 111.44	Weak
Bar-Oz, 2007	Cardiovascular malformations	First trimester paroxetine exposure in pregnancy	188/16501	6	Cohort, case-control	OR	1.79 (1.27, 2.53)	0.001	2.16 (1.25, 3.72)	0	0.79	1.10, 2.92	Weak
Bar-Oz, 2007	Major malformations	First trimester paroxetine exposure in pregnancy	646/16880	7	Cohort, case-control	OR	1.33 (1.03, 1.71)	0.03	1.83 (0.59, 5.64)	23	0.98	0.78, 2.26	Weak
Rahimi, 2006	Major malformations	SSRI during pregnancy,	97/2529	9	Cohort	OR	1.40 (0.86, 2.33)	0.17	1.12 (0.08, 17.78)	0	0.87	0.76, 2.34	No association
Rahimi, 2006	Cardiovascular malformations	SSRI during pregnancy,	21/1753	8	Cohort	OR	1.18 (0.36, 3.89)	0.78	3.40 (0.21, 251.89)	0	0.43	0.25, 5.65	No association
Rahimi, 2006	Minor malformations	SSRI during pregnancy	88/500	2	Cohort	OR	0.93 (0.13, 6.60)	0.94	0.39 (0.22, 0.71)	0	NA	NA	No association
Rahimi, 2006	Spontaneous abortion	SSRI during pregnancy	225/2378	5	Cohort	OR	1.70 (1.26, 2.30)	0.001	2.06 (0.86, 5.17)	0	0.92	1.04, 2.78	Weak
Healy, 2016	Autistic Spectrum or related disorders	SSRI during pregnancy	735/83750	5	Case-control	OR	1.99 (1.67, 2.36)	5.0x10 ⁻¹⁵	2.3 (1.23, 2.74)	0	0.57	1.50, 2.63	Weak
Oh, 2014	Risk of coronary heart disease	SSRI	84709/925749	12	Cohort, case-control	OR	0.93 (0.65, 1.33)	0.672	1.29 (0.89, 1.87)	98	0.400	0.24, 3.60	No association
McDonagh, 2014	Preterm birth	Antidepressant during pregnancy	NR / 304	2	Cohort, case-control	OR	1.84 (0.79, 4.27)	0.155	1.73 (0.63, 4.58)	0	NA	NA	No association
Huybrechts, 2014	Preterm birth	Antidepressant during early pregnancy (unadjusted)	NR/308762	8	Cohort, case-control	OR	1.58 (1.31, 1.92)	2.0x10 ⁻⁵	1.38 (0.91, 2.10)	14	0.22	1.12, 2.23	Weak

Huybrechts, 2014	Preterm birth	Antidepressant during pregnancy (any time unadjusted)	NR/1559757	4	Cohort, case-control	OR	1.44 (1.37, 1.51)	6.0×10^{-50}	1.44 (1.37, 1.51)	0	0.50	1.30, 1.61	Suggestive
Huybrechts, 2014	Preterm birth	Antidepressant during pregnancy (any time adjusted)	NR/294792	17	Cohort, case-control	OR	1.57 (1.41, 1.75)	$3. \times 10^{-16}$	1.21 (0.67, 2.21)	19	0.00	1.26, 1.97	Suggestive
Myles, 2013	Major malformations	Fluoxetine during first trimester	NR/NR	9	Cohort, case-control	OR	1.14 (1.01, 1.30)	0.04	0.79 (0.56, 1.12)	32	0.58	0.76, 1.78	Weak
Myles, 2013	Major malformations	Paroxetine during first trimester	NR/NR	8	Cohort, case-control	OR	1.29 (1.11, 1.49)	0.001	1.00 (0.06, 16.85)	0	0.28	1.07, 1.55	Weak
Myles, 2013	Major malformations	Sertraline during first trimester	NR/NR	6	Cohort, case-control	OR	1.01 (0.88, 1.17)	0.88	0.62 (0.09, 4.00)	0	0.84	0.83, 1.24	No association
Myles, 2013	Major malformations	SRI during first trimester	NR/NR	6	Cohort, case-control	OR	1.06 (0.93, 1.21)	0.37	0.97 (0.81, 1.16)	0	0.59	0.88, 1.28	No association
Myles, 2013	Cardiovascular malformations	Fluoxetine during first trimester	NR/NR	6	Cohort, case-control	OR	1.21 (0.99, 1.48)	0.07	0.77 (0.19, 3.11)	0	0.57	0.91, 1.61	No association
Myles, 2013	Cardiovascular malformations	Paroxetine during first trimester	NR/NR	8	Cohort, case-control	OR	1.44 (1.16, 1.79)	0.001	0.88 (0.21, 0.80)	6	0.88	1.03, 2.01	Weak
Myles, 2013	Cardiovascular malformations	Sertraline during first trimester	NR/NR	8	Cohort, case-control	OR	0.97 (0.64, 1.48)	0.90	0.65 (0.34, 1.25)	63	0.34	0.25, 3.87	No association
Myles, 2013	Minor malformations	SSRI during first trimester	NR/NR	7	Cohort, case-control	OR	1.18 (0.84, 1.66)	0.34	0.62 (0.20, 1.92)	32	0.33	0.54, 2.59	No association

Nikfar, 2012	Cardiovascular malformations	SSRI during pregnancy,	25128/ 2309472	19	Cohort	OR	1.17 (0.44, 3.09)	0.75	2.17 (0.92, 4.35)	98	0.41	0.02, 70.99	No association
Nikfar, 2012	Minor malformations	SSRI during pregnancy,	2435/ 698770	6	Cohort	OR	1.35 (0.61, 3.00)	0.45	1.06 (0.70, 1.56)	89	0.78	0.10, 18.31	No association
Biffi, 2017	Acute heart disease	TCAs	71838/ 741948	9	Cohort, case-control	RR	1.29 (1.09, 1.54)	0.00	1.41 (1.37, 1.45)	73	0.47	0.81, 2.07	Weak
Biffi, 2017	Haemorrhagic stroke	SSRI	1054/ 159014	3	Cohort, case-control	RR	1.33 (0.86, 2.06)	0.20	1.00 (0.61, 1.63)	38	0.43	0.02, 85.96	No association
Biffi, 2017	Ischaemic stroke	SSRI	4281/ 159014	3	Cohort, case-control	RR	1.15 (0.98, 1.36)	0.09	1.10 (0.88, 1.37)	0	0.37	0.39, 3.37	No association
Zhang, 2017	Atrial septal defect	SSRI during pregnancy	4096/ 2041138	6	Cohort	RR	2.06 (1.40, 3.03)	2.3×10^{-4}	2.60 (1.84, 3.68)	58	0.79	0.67, 6.28	Suggestive
Zhang, 2017	Ventricular septal defect	SSRI during pregnancy	12563/ 2990642	7	Cohort	RR	1.15 (0.97, 1.36)	0.11	1.20 (1.04, 1.39)	30	0.11	0.78, 1.68	No association
Ross, 2013	Spontaneous abortion	SSRI during pregnancy	367/ 5780	3	Cohort	RR	1.46 (0.99, 2.16)	0.06	1.48 (0.83, 2.66)	0	0.75	0.12, 18.58	No association
Gao, 2017	Major congenital malformations	Fluoxetine during pregnancy	132646/ 4234692	12	Cohort	RR	1.18 (1.08, 1.29)	3.5×10^{-4}	1.25 (1.10, 1.42)	0	0.49	1.06, 1.30	Suggestive
Gao, 2017	Cardiovascular malformations	Fluoxetine during pregnancy	64921/ 6385820	12	Cohort	RR	1.36 (1.17, 1.59)	9.0×10^{-5}	1.34 (1.10, 1.63)	23	0.12	0.99, 1.87	Suggestive
Morales, 2018	Autism spectrum disorders	Antidepressant during pregnancy	28105 / 2519942	11	Cohort, case-control	RR	1.53 (1.31, 1.78)	4.0×10^{-8}	2.17 (1.20, 3.93)	33	0.82	1.07, 2.19	Convincing
Morales, 2018	Attention deficit hyperactivity disorder	Antidepressant during pregnancy	NR/ NR	7	Cohort, case-control	RR	1.38 (1.13, 1.69)	0.002	0.97 (0.54, 1.73)	53	0.74	0.80, 2.23	Weak
Morales, 2018	Attention deficit hyperactivity disorder	Antidepressant during preconception period	NR/ NR	5	Cohort, case-control	RR	1.38 (1.14, 1.69)	0.001	1.50 (1.01, 2.22)	25	0.96	0.86, 2.24	Weak

Morales, 2018	Attention deficit hyperactivity disorder	Antidepressant during pregnancy	NR/ NR	3	Sibling study design	RR	0.88 (0.70, 1.12)	0.296	0.70 (0.37, 1.31)	0	0.08	0.19, 4.00	No association
Gao, 2018	Major congenital malformations	SSRI during pregnancy	NR/ NR	9	Cohort	RR	1.11 (1.03, 1.19)	0.004	1.13 (1.06, 1.20)	38	0.83	0.94, 1.31	Weak
Gao, 2018	Cardiovascular malformations	SSRI during pregnancy	NR/ NR	18	Cohort	RR	1.24 (1.11, 1.37)	6.2x10 ⁻⁵	1.03 (0.86, 1.24)	59	0.45	0.89, 1.71	Weak

TCAs– tricyclic antidepressants, SSRI – selective serotonin reuptake inhibitors, I^2 –heterogeneity, CI – confidence interval, OR – odds ratio, RR – relative risk, HR– hazard ratio, NA– not applicable.

eTable 7 List of covariates used for the sensitivity analysis limited to studies adjusted for covariates

Meta-analysis author, year, outcome	Primary study author year	Factors considered during adjusted analysis
Man, 2018 Attention deficit hyperactivity disorder	Boukhris, 2017	Gender, birth year, maternal age, maternal education level, recipient of social assistance, area of residence, maternal psychiatric disorders in the year prior to or during pregnancy (depression/anxiety, other psychiatric disorders), maternal comorbidities (gestational diabetes, gestational hypertension), maternal history of ADHD
	Laugesen, 2013	Gender of the child, calendar time at birth, birth order, maternal age at birth, maternal smoking status, maternal psychiatric diagnoses, paternal psychiatric diagnoses, maternal diseases during pregnancy (infections, epilepsy) and maternal anxiolytics/ hypnotics/sedatives use during pregnancy
	Malm, 2016	Sex; socioeconomic status; smoking during pregnancy; neonatal care unit; maternal history of other psychiatric diagnosis; maternal history of substance abuse; paternal history of psychiatric diagnosis; parental death
	Man, 2017	Maternal age at delivery, infant's sex, birth year, birth hospital, parity, maternal underlying medical conditions before delivery (pre-existing diabetes, epilepsy, gestational diabetes, psychiatric conditions, hypertension), use of other psychotropic drugs (antipsychotics, British National Formulary chapter 4.2.1, 4.2.2), and socioeconomic status.
	Sujan, 2017	Parity; year of birth; country of birth; age at childbearing; highest level of completed education; history of any criminal conviction; history of severe psychiatric illnesses (inpatient diagnosis of ICD-8, ICD-9, or ICD-10 schizophrenia, bipolar disorder, or other non-drug-induced psychoses); and history of any suicide attempts.
	Clements, 2015	Gender, race, birth year, insurance type, median income tertile, past history of maternal depression
	Figueroa, 2010	Maternal age group, gender of the child, urban or rural metropolitan statistical area, year of birth, age at last claim and at end of eligibility, maternal and paternal mental health diagnoses, the presence or absence of maternal mental health-related visits by period of time, the use of other psychotropics during pregnancy, and perinatal complications
Fu, 2018 Cataract risk	Becker, 2017	Calendar time (same index date), age, sex, general practice, and number of years of active history in the CPRD before the index date, BMI, smoking, diabetes, hypertension, and systemic steroids
	Chou, 2017	Age, sex, index date, patient's demographics, mental illness characteristics, propensity score derived from comorbid conditions, and concomitant medications
	Klein, 2001	Age and gender
Morales, 2018 Autism pre-pregnancy exposure	Boukhris, 2016	Gender, year of birth, maternal age, marital status, living alone, education, social assistance or care, maternal psychiatric history, paternal psychiatric history, maternal physical history, paternal physical history
	Brown, 2017	Gender, gestational age at delivery, maternal age, maternal psychiatric history, maternal physical history, pre-pregnancy related/delivery, severity of depression, parity, drugs other than antidepressants
	Castro, 2016	Gender, year of birth, gestational age at delivery, maternal age, education, maternal psychiatric history, maternal physical history, severity of depression, parity, insurance type ethnicity or country of origin, maternal income

	Clements, 2015	Gender, year of birth, birth weight, gestational age at delivery, maternal age, education, maternal psychiatric history, maternal physical history, pre-pregnancy related/delivery, severity of depression, parity, insurance type ethnicity or country of origin
	Croen, 2011	Gender, year of birth, birth weight, gestational age at delivery, maternal age, education, maternal psychiatric history, parity, ethnicity or country of origin
	Hviid, 2013	Gender, year of birth, gestational age at delivery, maternal age, education, maternal psychiatric history, smoking status, parity, ethnicity or country of origin residence, employment status
	Sujan, 2017	Year of birth, maternal age, education, maternal psychiatric history, parity, ethnicity or country of origin,
Laporte, 2017 Severe bleeding at any site	de Abajo 1999	UGIB history, smoking status, current use of NSAID, AC, corticosteroids, aspirin
	de Abajo 2000	Hypertension, migraine, asthma or COPD, smoking status, BMI, current use of NSAIDs
	Bak 2002	Age, sex, hypertension, diabetes, smoking status, AC, antiarrhythmics, antianginal drugs
	Meijier 2004	Bleeding history, NSAIDs, AC, glucocorticoids, estrogens, progestones, histamine blockers, PPIs, antidiabetic agents
	Kurdyak 2005	UGIB history, current use of aspirin, NSAID, glucocorticoid, PPIs, H2 reuptake inhibitors
	Tata 2005	NR
	Helin-Salmivaara 2007	Histamine-2 receptor antagonist, plain misoprostol, PPIs, warfarin, clopidogrel or inhaled glucocorticoid and tramadol, hospitalisation for arthroplasty, hypertension, angina pectoris, cardiac insufficiency, diabetes mellitus, rheumatoid arthritis, asthma
	Kharof 2007	NR
	Vonbach 2007	Glucocorticoids, NSAIDs, AC, SSRIs, TAI, PPIs, hypertensive disease
	Ziegelstein 2007	NR
	de Abajo 2008	Age, sex, calendar year, smoking status, alcohol intake, history of GI disorder, NSAIDs, systemic corticosteroids, warfarin, low-dose aspirin, antiplatelet drug
	Lewis 2008	Age, sex, race, alcohol consumption, history of ulcer disease, hypertension, PPI use, H2RA use, ASA dose and NSAID dose
	Opatrny 2008	Age, sex, BMI, blood pressure, smoking status, comorbid conditions, warfarin, clopidogrel, antidepressant
	Salkeld 2008	Previous PPH, multiple pregnancy, prolonged labor, abnormalities of the forces of labor, obstructed labor, perineal laceration or other gynecologic laceration, other obstetric trauma, placenta previa, placental abruption, and hypertensive disorders of pregnancy
	Schalekamp 2008	NR
	Vidal 2008	History of peptic ulcer, dyspepsia, UGIB, diabetes mellitus, smoking habit, alcohol consumption and use of antacids, PPIs, sucralfate, nitrates, systemic NSAIDs, topical NSAIDs, analgesics, antiplatelet drugs, dihydropyridine calcium antagonists and statin
	Barbui 2009	Age, gender, use of antianemic preparations, use of drugs for peptic ulcer

	Chen 2009	Use of aspirin, AC, risperidone, anxiety, alcohol abuse, substance abuse, hypertension, diabetes, hypercholesterolemia, cardiac diseases
	Dall 2009	Age, gender, calendar year, low dose aspirin, PPIs, NSAIDs, alcohol abuse, cerebral ischemia, stroke, warfarin, clopidogrel, dipyridamol, steroids, helicobacter eradication, peptic ulcer, UGIB, cirrhosis
	Targownik 2009	Cardiovascular disease, respiratory disease, hepatic disease, renal disease, active malignancy, alcohol abuse, depression, schizophrenia, acute hospitalisation, upper endoscopy, H2-receptor antagonists, warfarin, clopidogrel, systemic corticosteroids, tricyclic antidepressants
	Carvajal 2011	Alcohol and caffeine consumption, past history of GI disorders, family history of GI bleeding, osteoarthritis, number of medicines taken and use of NSAIDs, salicylates, PPIs, H2 antihistamines, antacids, antiplatelet agents and AC
	Douglas 2011	Smoking, alcohol, BMI, prior history of transient ischemic attack or other stroke, hypertension, diabetes, NSAID use, aspirin use, clopidogrel or dipyridamole use, year of first prescription, observation time
	Verdel 2011	NSAIDs, oral glucocorticoids, PPI, platelet aggregation inhibitors
	de Abajo 2013	Age, gender, calendar year, smoking, peptic ulcer history, number of GP visits in the year prior to index date and concomitant use of other medications
	Andreasen 2006	Age, sex, preoperative use of platelet inhibitors, NSAIDs, oral anticoagulant, place of surgery, use of extracorporeal circulation, concomitant valve surgery, Charlson comorbidity index
	Hauta-Aho 2009	Age, sex, study ward, PPI and oral glucocorticoid medications
	Kim 2009	NR
	Gärtner 2010	Age
	Tully 2012	Propensity score (including age, sex, urgency of surgery, previous myocardial infarction, respiratory disease, left ventricular ejection fraction, diabetes mellitus, renal disease, peripheral vascular disease, cerebrovascular disease, cardiogenic shock, heart failure, hypertension, smoking and OPCAB procedure, statin, antiplatelet, anticoagulants)
	Basile 2013	Age, body weight, surgery type
	Mortensen 2013	Propensity-matched analysis (among 5837 users and 30338 non-users) and use of other drugs during follow-up (non-SSRI, other antidepressant, blood pressure lowering drugs, platelet inhibitors, VKA, statins)
	Seitz 2013	Age, sex, Charlson score, number of medications, residence (long-term care or community)
	Quinn 2013	Time varying ATRIA bleeding risk score, INR value
Jiang, 2016 Postpartum hemorrhage	Salkeld 2008	Previous postpartum hemorrhage, multiple pregnancy, prolonged labor, abnormalities of the forces of labor, obstructed labor, perineal laceration or other gynecologic laceration, other obstetric trauma, placenta previa, placental abruption, and hypertensive disorders of pregnancy
	Palmsten 2013	Delivery year, age, race, multiple pregnancy, diabetes, coagulopathy, number of outpatient mood/anxiety disorder diagnoses, number of inpatient mood/anxiety disorder diagnoses, psychotic disorder, other mental health disorder, pain indication, sleep disorder, anticonvulsant dispensing, benzodiazepine dispensing, aspirin dispensing, heparin dispensing, low molecular weight heparin dispensing, warfarin dispensing, and number of outpatient visits and days in hospital during baseline.
	Lindqvist 2014	Maternal age, parity, BMI, educational level, smoking, coagulation defects, history of previous abortion/miscarriage, placental abruption, placenta previa, and maternal depressive symptoms.

	Lupattelli 2014	Maternal age, marital status, BMI, smoking, placenta previa, bleeding episode in first trimester, and depressive symptoms.
	Grzeskowiak 2015	Delivery year, age, socio-economic status, race, multiple pregnancy, parity, smoking status, alcohol or substance abuse during pregnancy, coagulation defects, asthma, diabetes, hypertension, previous caesarean section, and use of other psychotropic medications
	Joseph 2015	Social assistance, residence urban, previous caesarean, multi-foetal pregnancy, placenta previa/ abruption, polyhydramnios, prolonged labour, preeclampsia/eclampsia, epidural analgesia, labour induction, uterine rupture, cervical laceration, caesarean delivery, perineal laceration, vaginal laceration, chorioamnionitis.
	Hanley 2016	Year of birth, maternal age, parity, preterm birth, multifetal pregnancy, diabetes (both gestational and pre-existing), coagulopathy, smoking during pregnancy, blood thinner use in the month before delivery, anxiolytic use in the month before delivery, antipsychotic use in the month before delivery, a diagnosis of mood disorder, any psychiatric visits, or any psychiatric hospitalization in the 5 months before delivery.
	Kim 2016	Age, race, education, parity, comorbid anxiety to the SRI exposure variable, and depressive symptoms.
Jiang, 2015 Upper gastrointestinal bleeding	de Abajo 1999	UGIB history, smoking status, current use of NSAID, AC, corticosteroids, aspirin
	Helin-Salmivaara 2007	Histamine-2 receptor antagonist, plain misoprostol, PPIs, warfarin, clopidogrel or inhaled glucocorticoid and tramadol, hospitalisation for arthroplasty, hypertension, angina pectoris, cardiac insufficiency, diabetes mellitus, rheumatoid arthritis, asthma
	de Abajo 2008	Age, sex, calendar year, smoking status, alcohol intake, history of GI disorder, NSAIDs, systemic corticosteroids, warfarin, low-dose aspirin, antiplatelet drug
	Schalekamp 2008	NR
	Opatrny 2008	Age, sex, BMI, blood pressure, smoking status, comorbid conditions, warfarin, clopidogrel, antidepressant
	Lewis 2008	Age, sex, race, alcohol consumption, history of ulcer disease, hypertension, PPI use, H2RA use, ASA dose and NSAID dose
	Vidal 2008	History of peptic ulcer, dyspepsia, UGIB, diabetes mellitus, smoking habit, alcohol consumption and use of antacids, PPIs, sucralfate, nitrates, systemic NSAIDs, topical NSAIDs, analgesics, antiplatelet drugs, dihydropyridine calcium antagonists and statin
	Barbui 2009	Age, gender, use of antianemic preparations, use of drugs for peptic ulcer
	Targownik 2009	Cardiovascular disease, respiratory disease, hepatic disease, renal disease, active malignancy, alcohol abuse, depression, schizophrenia, acute hospitalisation, upper endoscopy, H2-receptor antagonists, warfarin, clopidogrel, systemic corticosteroids, tricyclic antidepressants
	Dall 2009	Age, gender, calendar year, low dose aspirin, PPIs, NSAIDs, alcohol abuse, cerebral ischemia, stroke, warfarin, clopidogrel, dipyridamol, steroids, helicobacter eradication, peptic ulcer, UGIB, cirrhosis
	Carvajal 2011	Alcohol and caffeine consumption, past history of GI disorders, family history of GI bleeding, osteoarthritis, number of medicine taken and use of NSAIDs, salicylates, PPIs, H2 antihistamines, antacids, antiplatelet agents and AC
	Verdel 2011	NSAIDs, oral glucocorticoids, PPI, platelet aggregation inhibitors
	de Abajo 2013	Age, gender, calendar year, smoking, peptic ulcer history, number of GP visits in the year prior to index date and concomitant use of others medications

	Wang 2014	Concurrent drug use
	Huang 2011	Age, concurrent drug use
Huang, 2014 Preterm birth	Nordeng 2012	Depression
	Yonkers 2012	Depression
	Latendresse 2011	Depression
	Lewis 2010	Depression
	Lenneval 2007	NR
	Suri 2007	Depression
Wu, 2013 Fracture risk, TCAs	Ensrud 2003	Age, health status, use of ≥ 1 medication, walking for exercise, functional impairment, fall in previous year, cognitive function, weight change, gait speed, inability to rise from chair, femoral neck BMD
	Lewis 2007	Age, BMD
	Ziere 2008	Age, sex, depression during follow-up period, disability category, lower-limb disability
	Diem 2011	Age, health status, IADL, ability to rise from chair, m-MMSE, smoking, alcohol use, estrogen use, bisphosphonate use, benzodiazepine use, thiazide use, PPI use, oral steroid use, weight, GDS score, walks for exercise, history of prior fracture, total-hip BMD and history of falls in previous year
	Coupland 2011	Age, sex, depression, deprivation, smoking status, comorbidities (ischemic heart disease, DM, HTN, stroke, cancer, dementia, epilepsy or seizures, PD, hypothyroidism, OCD) and use of other drugs (eg, statins, NSAIDs, antipsychotics, lithium, aspirin, anti-HTN drugs, anticonvulsants, hypnotics, anxiolytics)
	Ray 1987	Sex, race, age, index year, and home status, diagnosis of dementia
	Ray 1991	Age, sex, calendar year, nursing home residence on index date, and for hospitalization and use of specific medications (narcotic analgesics, anti-HTN, and other cardiovascular drugs) in year preceding index date
	Liu 1998	Age, sex, comorbidity (eg, depression, dementia, osteoporosis), previous drug exposure (eg, sedative, tranquilizer, cardiac drug, anti-PD agent, thyroid-replacement drug, anticonvulsant, insulin, glucocorticoid, estrogen, etidronate)
	Hubbard 2003	Age, sex, general practice, duration of available GPRD data, history of falls, and history of prescriptions for hypnotics and antipsychotics
	Vestergaard 2006	Age, sex, psychiatric comorbidity (eg, manic depression, schizophrenia, alcoholism, eating disorder); medication use (eg, anxiolytic, sedative, neuroleptic, corticosteroid, antiepileptic, cardiovascular agent, lithium); hospital stay; prior fracture; income; working, educational, and residential status; Charlson index
	Van den Brand	Age, sex, geographical region, other antidepressant, use of benzodiazepine, antipsychotic, lithium, anti-PD agent, anticonvulsant, oral-inhaled corticosteroid, bronchodilator, HRT, antiarrhythmic, thiazide diuretic, β -blocker, opioids, anticonvulsants, DM drug, ≥ 2 dispensings of an NSAID, DMARDs, metoclopramide hydrochloride; history of malignant neoplasm, mental disorder, cerebrovascular disease, obstructive airway disease, or IBD
	Verdel 2010	Age; sex; geographical area; calendar time; cancer; cardiovascular disease; cerebrovascular disease; IBD; mental disorder; obstructive airway disease; use of antidiabetic, antiepileptic, anti-PD drug, antipsychotic, benzodiazepine, β -blocker, DMARD, HRT, NSAID, oral glucocorticoid, opioid

Ross, 2013 Apgar score at 5 minutes	Simon et al., 2002	maternal age, year of delivery, lifetime number of antidepressant prescriptions filled or refilled, lifetime history of outpatient psychiatric treatment, lifetime history of inpatient psychiatric treatment, tobacco use, other substance use, race, number of prior births
	Laine et al., 2003	maternal age, gravidity, parity, gestational age, time and mode of delivery
	Zeskind & Stephens, 2004	maternal age (± 2 yrs), maternal cigarette use, low socioeconomic status,
	Pearson et al., 2007	age (± 5 years), parity, tobacco use, marital status
	Calderon-Margalit et al., 2009	maternal age, race, marital status, education, smoking during pregnancy, preeclampsia, parity, singleton pregnancy
	Lund et al., 2009	maternal age, BMI, smoking, a previous pregnancy with prematurity, gestational age, previous birth of a low-birth-weight infant, parity, and coffee and alcohol intake
	Wisner et al., 2009	maternal age, race
	Reis et al., 2010	year of birth, maternal age, parity, smoking, body mass index (BMI)
Oderda, 2012 Hip fracture	van den Brand (2009)	Age sex, medications, comorbidities
	Chang (2008)	Age, sex
	Ensrud (2003)	Age, debilitation, gait speed, femoral neck, bone density
	Hubbard (2003)	History of falls, medications
	Wang (2001)	Age, sex, race, medications, debilitation
	Jacqmin-Gadda (1998)	Age, sex, residence
	Guo (1998)	Comorbidities, medications
	Liu (1998)	Comorbidities, medications
	Johansson (1996)	Sex, comorbidities
	Lichtenstein (1994)	Age, sex, comorbidities
	Cumming (1993)	Age, sex, residence
	Jensen (1991)	Age, sex, debilitation, medications
	Ray (1991)	Age, sex, debilitation, medications
	Ray (1987)	Age, sex, race, residence
Barbui, 2009 Suicide attempt and completion	Olfson 2006	White population, median income, number of per capita physicians pediatricians, child psychiatrists and psychiatrists
	Olfson 2006	Age, sex, race, substance use disorder, recent suicide attempt, treatment with antipsychotic anxiolytic, hypnotic, stimulant and mood stabilizing drugs
	Sondergard 2007	Age, sex, and socioeconomic classification

	Tiihonen 2006	Age, sex, geographical location, number of suicide attempts before the index hospitalization, number of suicide attempts during follow-up, use of multiple antidepressants and number of purchased antidepressants during the previous year
	Gibbons 2007	No
	Valuck 2004	Propensity score and calendar year
Khanassov, 2018 Fracture risk; SSRIs	van den Brand, 2009	Age, sex, geographical region, other than SSRIs antidepressants, benzodiazepines, antipsychotics, lithium, anti-Parkinson drugs, anticonvulsants, corticosteroids, hormone-replacement therapy, disease-modifying antirheumatic drugs, nonsteroidal antiinflammatory drugs, antiarrhythmic, thiazide diuretics, beta-blockers, opiates, metoclopramide, antidiabetic drugs; history of hospitalization
	Verdel, 2010	Age, sex, geographical region, calendar time, other than SSRIs antidepressants, benzodiazepines, antipsychotics, lithium, anti-Parkinson drugs, anticonvulsants, corticosteroids, hormone-replacement therapy, disease-modifying antirheumatic drugs, nonsteroidal antiinflammatory drugs, antiarrhythmic, thiazide diuretics, beta-blockers, opiates, metoclopramide, antidiabetic drugs, thyroid hormones; history of hospitalization
	Vestergaard, 2006	Age, sex, psychiatric comorbidity (eg, schizophrenia, alcoholism), medication use such as anxiolytic, sedative, neuroleptic, corticosteroid, antiepileptic, lithium, hospitalization, prior fracture, income, working status, education, residence, Charlson index
	Bolton, 2008	Age, sex, ethnicity, income, residence, comorbidity index of the John Hopkins Ambulatory Care Group system (diabetes, ischemic heart disease, myocardial infarction, hypertension, epilepsy, rheumatoid arthritis, organ transplantation, COPD, home care use, depression, substance abuse, dementia, schizophrenia), medication use such as anticonvulsants, diuretics, anticoagulant, thyroid hormone
	Liu, 1998	Age, sex, comorbidities (depression, dementia and other), medications such as sedatives, tranquilizers, cardiac agents, anti-Parkinson drugs, thyroid hormones, anticonvulsants, diabetic agents, corticosteroids, estrogens, etidronate; different exposure categories and doses of antidepressant
	Abrahamsen, 2009	Age, previous fracture, modified Charlson comorbidity index, groups of medications
	Wang, 2016	Age, sex, hypertension, diabetes, osteoporosis, history of falls, cardiac diseases, chronic obstructive pulmonary disease, urinary incontinence, Parkinson disease, chronic mental disorders, dementia, depression, liver disease, peripheral vascular disease, cerebrovascular disease, arthritis, chronic kidney diseases, glaucoma; use of medications (opiates, nonopioid analgesics, antipsychotics, anxiolytics, sedatives, corticosteroids, hormone replacement therapy, antiepileptics, tricyclics)
	Rabenda, 2012	Lumbar spine, femoral neck and total hip BMD, depression, history of vertebral/ nonvertebral fracture, benzodiazepine use, antihypertensive and antiarrhythmic drugs
	Spangler, 2008	Age, BMD, height, weight, ethnicity, physical function, hormone replacement therapy, smoking, years since menopause, number of falls, previous fracture, use of analgesics, narcotics, cardiovascular diseases
	Richards, 2007	Age, sex, education, study center, BMI, comorbidities based on modified Charlson index, self-reported general health, prior smoking and alcohol intake, history of falls and fractures, calcium and vitamin D intake, dementia based on MMSE, depression, medication use such as bisphosphonates, antihypertensive, diuretics, corticosteroids, estrogens, benzodiazepines, anticonvulsants, antipsychotics, tricyclics, BMD
	Lewis, 2007	Age, BMD
	Schneeweiss, 2004	BMI, current smoking status, activities of daily living, cognitive impairment

	Carriere, 2016	Age, center, sex, smoking, benzodiazepines, other CNS drugs, osteo-articular pain, time since first depressive episode, antiosteoporosis drugs, corticosteroids
	Ensrud, 2003	Age, health status, one or more medical conditions, walking for exercise, functional impairment, fall in previous year, cognitive function, weight change, gait speed, inability to rise from chair, femoral neck bone density.
	Ziere, 2008	Age, sex, disability category, lower limb disability, depression
	Diem, 2011	Age, health status, instrumental daily activities, ability to rise from chair, MMSE, smoking, alcohol use, hormone replacement therapy, bisphosphonate use, benzodiazepine use, thiazide use, proton pump inhibitor use, corticosteroid use, weight, depression, walks for exercise, prior fracture, BMD
	Coupland, 2011	Age, sex, year of depression diagnosis, diagnosis of depression before age 65, severity of index depression, deprivation level, smoking status, comorbidities (heart conditions, diabetes, dementia, cancer, epilepsy, Parkinson's disease, hypothyroidism, obsessive-compulsive disorder, stroke), use of statins, nonsteroidal antiinflammatory drugs, antipsychotics, lithium, aspirin, antihypertensive, anticonvulsants, hypnotics), previous falls.
	Bakken, 2013	Sex, birth year, time period
	Cheng, 2016	Age, sex, urbanization, osteoporosis, Charlson comorbidity index
	Souverein, 2016	Age, sex, previous fracture, corticosteroids, rheumatoid arthritis, smoking, alcohol use, BMI, osteoporosis, history of bone diseases, previous use of bisphosphonates or other bone protecting drugs: raloxifene, strontium ranelate, parathyroid hormone, calcium, vitamin D, calcitonin, calcitriol
	Sheu 2015	Age, sex, previous fracture, corticosteroids, rheumatoid arthritis, smoking, alcohol use, BMI, osteoporosis, history of bone diseases, previous use of bisphosphonates or other bone protecting drugs: raloxifene, strontium ranelate, parathyroid hormone, calcium, vitamin D, calcitonin, calcitriol
	Adachi,2014	Age, BMI, parental history of hip fracture, rheumatoid arthritis, prior fracture, osteoarthritis, celiac disease, Crohn's disease, Parkinson's disease, falls in the past year, smoking, alcohol intake, anxiety, depression, general health, physical function and vitality

PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	3–4
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	6
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	7
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	7
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	7-8
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	7
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	eBox1
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	8 Supplement methods
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	9
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	9

Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	9
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	9
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	9-10

Page 1 of 2

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	11
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	12-13
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	Figure 1 eTable 1
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	Table 1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Table 1
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Tables 2-3, eTables 2-4
Synthesis of results	21	Present the main results of the review. If meta-analyses are done, include for each, confidence intervals and measures of consistency.	12-13
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	Tables 2-3, eTables 2-4
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	13 Supplement results
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	13-16

Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	16-17
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	17
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	18-19

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

MOOSE Checklist for Meta-analyses of Observational Studies

Item No	Recommendation	Reported on Page No
Reporting of background should include		
1	Problem definition	6-7
2	Hypothesis statement	7
3	Description of study outcome(s)	7-8
4	Type of exposure or intervention used	7-8
5	Type of study designs used	7-8
6	Study population	7-9
Reporting of search strategy should include		
7	Qualifications of searchers (eg, librarians and investigators)	NA
8	Search strategy, including time period included in the synthesis and key words	7, eBox1
9	Effort to include all available studies, including contact with authors	7
10	Databases and registries searched	7
11	Search software used, name and version, including special features used (eg, explosion)	NA
12	Use of hand searching (eg, reference lists of obtained articles)	7
13	List of citations located and those excluded, including justification	Figure 1, eTable 1
14	Method of addressing articles published in languages other than English	We placed restrictions on English.
15	Method of handling abstracts and unpublished studies	NA
16	Description of any contact with authors	NA
Reporting of methods should include		
17	Description of relevance or appropriateness of studies assembled for assessing the hypothesis to be tested	7-8
18	Rationale for the selection and coding of data (eg, sound clinical principles or convenience)	8-9
19	Documentation of how data were classified and coded (eg, multiple raters, blinding and interrater reliability)	8-9
20	Assessment of confounding (eg, comparability of cases and controls in studies where appropriate)	9-10
21	Assessment of study quality, including blinding of quality assessors, stratification or regression on possible predictors of study results	9-10
22	Assessment of heterogeneity	9-10
23	Description of statistical methods (eg, complete description of fixed or random effects models, justification of whether the chosen models account for predictors of study results, dose-response models, or cumulative meta-analysis) in sufficient detail to be replicated	9-10
24	Provision of appropriate tables and graphics	Tables 1-4, Figure 1, Supplement file
Reporting of results should include		

25	Graphic summarizing individual study estimates and overall estimate	Tables 2-3, Supplement file
26	Table giving descriptive information for each study included	Table 1
27	Results of sensitivity testing (eg, subgroup analysis)	Table 4, Supplement file
28	Indication of statistical uncertainty of findings	Supplement file

Item No	Recommendation	Reported on Page No
Reporting of discussion should include		
29	Quantitative assessment of bias (eg, publication bias)	NA
30	Justification for exclusion (eg, exclusion of non-English language citations)	NA
31	Assessment of quality of included studies	11, Table 4
Reporting of conclusions should include		
32	Consideration of alternative explanations for observed results	13-17
33	Generalization of the conclusions (ie, appropriate for the data presented and within the domain of the literature review)	14-17
34	Guidelines for future research	17
35	Disclosure of funding source	18-19

From: Stroup DF, Berlin JA, Morton SC, et al, for the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) Group. Meta-analysis of Observational Studies in Epidemiology. A Proposal for Reporting. *JAMA*. 2000;283(15):2008-2012. doi: 10.1001/jama.283.15.2008.

Transcribed from the original paper within the NEUROSURGERY® Editorial Office, Atlanta, GA, United States. August 2012.

Supplementary references

1. Ng QXV, N.; Ho, C. Y. X.; Sim, W. S.; Lim, D. Y.; Yeo, W. S. Selective Serotonin Reuptake Inhibitors and Persistent Pulmonary Hypertension of the Newborn: An Update Meta-Analysis. *Journal of women's health* (2002). 2019;28(3):331-338.
2. Patel KRI, S.; Singam, V.; Rastogi, S.; Silverberg, J. I. Association between atopic dermatitis, depression, and suicidal ideation: A systematic review and meta-analysis. *Journal of the American Academy of Dermatology*. 2019;80(2):402-410.
3. Rexwinkel RZ, J.; van Etten-Jamaludin, F. S.; Benninga, M. A.; Tabbers, M. M. Side effects associated with pharmacotherapy for pediatric irritable bowel syndrome and functional abdominal pain - not otherwise specified: a systematic review. *Expert opinion on drug safety*. 2019;18(2):111-125.
4. Perlman KB, D.; Israel, S.; Rollins, C.; Brown, E.; Tunteng, J. F.; You, R.; You, E.; Tanguay-Sela, M.; Snook, E.; Miresco, M.; Berlim, M. T. A systematic meta-review of predictors of antidepressant treatment outcome in major depressive disorder. *Journal of affective disorders*. 2019;243:503-515.
5. Amare ATS, K. O.; Tekola-Ayele, F.; Hsu, Y. H.; The association of obesity and coronary artery disease genes with response to SSRIs treatment in major depression. *Journal of neural transmission (Vienna, Austria : 1996)*. 2019;126(1):35-45.
6. Aguiar JPB, A. M.; Martins, A. P.; Leufkens, H. G. M.; Alves da Costa, F. Potentially inappropriate medications with risk of cardiovascular adverse events in the elderly: A systematic review of tools addressing inappropriate prescribing. *Journal of clinical pharmacy and therapeutics*. 2019.
7. Zhou C, Fang L, Chen Y, Zhong J, Wang H, Xie P. Effect of selective serotonin reuptake inhibitors on bone mineral density: a systematic review and meta-analysis. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2018;29(6):1243-1251.
8. Wu QL, B.; Tonmoy, S. Depression and risk of fracture and bone loss: an updated meta-analysis of prospective studies. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2018.
9. Wang SMH, C.; Bahk, W. M.; Lee, S. J.; Patkar, A. A.; Masand, P. S.; Pae, C. U. Addressing the Side Effects of Contemporary Antidepressant Drugs: A Comprehensive Review. *Chonnam medical journal*. 2018;54(2):101-112.
10. Visco DBM-d-C, R.; Chaves, W. F.; Lacerda, D. C.; Pereira, S. D. C.; Ferraz-Pereira, K. N.; Toscano, A. E. Selective serotonin reuptake inhibitors affect structure, function and metabolism of skeletal muscle: A systematic review. *Pharmacological research*. 2018;136:194-204.
11. Uguz F. Maternal antidepressant use during pregnancy and the risk of attention-deficit/hyperactivity disorder in children: A systematic review of the current literature. *Journal of clinical psychopharmacology*. 2018;38(3):254-259.
12. Tan ECKL, D.; Sandborgh-Englund, G.; Haasum, Y.; Johnell, K. Medications That Cause Dry Mouth As an Adverse Effect in Older People: A Systematic Review and Metaanalysis. *Journal of the American Geriatrics Society*. 2018;66(1):76-84.
13. Telang SW, C.; Olten, B.; Bloch, M. H. Meta-analysis: Second generation antidepressants and headache. *Journal of affective disorders*. 2018;236:60-68.
14. Sepehrpour AHE, M.; Jiskani, A.; Casula, R.; Athanasiou, T. Selective serotonin reuptake inhibitor use and outcomes following cardiac surgery-a systematic review. *Journal of thoracic disease*. 2018;10(2):1112-1120.
15. Prady SLH, I.; Fraser, L. K.; Mikocka-Walus, A. A systematic review of maternal antidepressant use in pregnancy and short- and long-term offspring's outcomes. *Archives of women's mental health*. 2018;21(2):127-140.
16. Cappetta KB, C.; Johnson, J. A.; Bloch, M. H. Meta-analysis: Risk of dry mouth with second generation antidepressants. *Progress in neuro-psychopharmacology & biological psychiatry*. 2018;84(Pt A):282-293.
17. Steinert TF, W. Epileptic Seizures Under Antidepressive Drug Treatment: Systematic Review. *Pharmacopsychiatry*. 2018;51(4):121-135.

18. Melo GD, K. L.; Rodrigues Filho, R.; Ortega, A. O. L.; Porporatti, A. L.; Dick, B.; Flores-Mir, C.; De Luca Canto, G. Association between psychotropic medications and presence of sleep bruxism: A systematic review. *Journal of oral rehabilitation*. 2018;45(7):545-554.
19. Moncrieff J. What does the latest meta-analysis really tell us about antidepressants? *Epidemiology and psychiatric sciences*. 2018;27(5):430-432.
20. Douros AA, M.; Renoux, C. Risk of Intracranial Hemorrhage Associated with the Use of Antidepressants Inhibiting Serotonin Reuptake: A Systematic Review. *CNS drugs*. 2018;32(4):321-334.
21. Chen LWC, M. Y.; Lian, Z. P.; Lin, H. S.; Chien, C. C.; Yin, H. L.; Chu, Y. H.; Chen, K. Y. Amitriptyline and Sexual Function: A Systematic Review Updated for Sexual Health Practice. *American journal of men's health*. 2018;12(2):370-379.
22. Comoretto RIR, F.; Lucenteforte, E.; Mugelli, A.; Trifiro, G.; Cascini, S.; Roberto, G.; Chinellato, A.; Filippelli, A.; Corrao, G. Bleeding events attributable to concurrent use of warfarin and other medications in high-risk elderly: meta-analysis and Italian population-based investigation. *European journal of clinical pharmacology*. 2018.
23. Varney AW, K.; Agius, M. What are the risks associated with the use of NSAIDs as an adjunct to SSRIs for treatment of depression? An evaluation of current evidence. *Psychiatry Danubina*. 2017;29(Suppl 3):375-382.
24. Uguz F. Is There Any Association Between Use of Antidepressants and Preeclampsia or Gestational Hypertension?: A Systematic Review of Current Studies. *Journal of clinical psychopharmacology*. 2017;37(1):72-77.
25. Salvi VG, I.; Cerveri, G.; Mencacci, C.; Barone-Adesi, F. The risk of new-onset diabetes in antidepressant users - A systematic review and meta-analysis. *PloS one*. 2017;12(7):e0182088.
26. Stubbs CM, L.; Crawford, S. A.; Wickersham, E. A.; Brockhaus, J. L.; McCarthy, L. H. Do SSRIs and SNRIs reduce the frequency and/or severity of hot flashes in menopausal women. *The Journal of the Oklahoma State Medical Association*. 2017;110(5):272-274.
27. Riediger CS, T.; Barlinn, K.; Maier, S.; Weitz, J.; Siepmann, T. Adverse Effects of Antidepressants for Chronic Pain: A Systematic Review and Meta-analysis. *Frontiers in neurology*. 2017;8:307.
28. Querido AL. Diving and antidepressants. *Diving Hyperb Med*. 2017;47(4):253-256.
29. Mezzacappa AL, P. A.; Gianfagna, F.; Cazas, O.; Hardy, P.; Falissard, B.; Sutter-Dallay, A. L.; Gressier, F. Risk for Autism Spectrum Disorders According to Period of Prenatal Antidepressant Exposure: A Systematic Review and Meta-analysis. *JAMA pediatrics*. 2017;171(6):555-563.
30. Maund EG, L. S.; Gøtzsche, P. C. Considering benefits and harms of duloxetine for treatment of stress urinary incontinence: A meta-analysis of clinical study reports. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne*. 2017;189(5):E194-E203.
31. Moraros JN, C.; Patten, S. B.; Mousseau, D. D. The association of antidepressant drug usage with cognitive impairment or dementia, including Alzheimer disease: A systematic review and meta-analysis. *Depression and anxiety*. 2017;34(3):217-226.
32. Locher CK, H.; Zion, S. R.; Werner, C.; Pine, D. S.; Kirsch, I.; Kessler, R. C.; Kossowsky, J. Efficacy and Safety of Selective Serotonin Reuptake Inhibitors, Serotonin-Norepinephrine Reuptake Inhibitors, and Placebo for Common Psychiatric Disorders Among Children and Adolescents: A Systematic Review and Meta-analysis. *JAMA psychiatry*. 2017;74(10):1011-1020.
33. Laux GB, B.; Hajak, G.; Lemke, M.; Volz, H. P. Pooled Analysis of Four Non-Interventional Studies: Effectiveness and Tolerability of the Antidepressant Agomelatine in Daily Practice. *Advances in therapy*. 2017;34(4):895-914.
34. Kaplan YCK-A, E.; Acar, S.; Sozmen, K. Maternal SSRI discontinuation, use, psychiatric disorder and the risk of autism in children: a meta-analysis of cohort studies. *British journal of clinical pharmacology*. 2017;83(12):2798-2806.
35. Heller HMR, A. C. J.; Bruning, A. H. L.; de Groot, C. J. M.; Scheele, F.; van Pampus, M. G.; Honig, A. Increased postpartum haemorrhage, the possible relation with serotonergic and other psychopharmacological drugs: a matched cohort study. *BMC pregnancy and childbirth*. 2017;17(1):166.

36. Hill LLL, V. L.; Winbrock, E. L.; Li, G.; Chihuri, S.; Lee, K. C. Depression, antidepressants and driving safety. *Injury epidemiology*. 2017;4(1):10.
37. Deidda AP, C.; Micheletto, L.; Bocchetta, A.; Del Zompo, M.; Stochino, M. E. Interstitial lung disease induced by fluoxetine: Systematic review of literature and analysis of Vigibase, Eudravigilance and a national pharmacovigilance database. *Pharmacological research*. 2017;120:294-301.
38. Brunnauer AL, G. Driving Under the Influence of Antidepressants: A Systematic Review and Update of the Evidence of Experimental and Controlled Clinical Studies. *Pharmacopsychiatry*. 2017;50(5):173-181.
39. Brown HKH-S, N.; Lunskey, Y.; Dennis, C. E.; Vigod, S. N. The Association Between Antenatal Exposure to Selective Serotonin Reuptake Inhibitors and Autism: A Systematic Review and Meta-Analysis. *The Journal of clinical psychiatry*. 2017;78(1):e48-e58.
40. Beyer CC, K.; Johnson, J. A.; Bloch, M. H. Meta-analysis: Risk of hyperhidrosis with second-generation antidepressants. *Depression and anxiety*. 2017;34(12):1134-1146.
41. Andrade C. Antidepressant Exposure During Pregnancy and Risk of Autism in the Offspring, 1: Meta-Review of Meta-Analyses. *The Journal of clinical psychiatry*. 2017;78(8):e1047-e1051.
42. Allain NL, C.; Falissard, B.; Allain, J. S.; Batail, J. M.; Polard, E.; Montastruc, F.; Drapier, D.; Naudet, F. Manic switches induced by antidepressants: an umbrella review comparing randomized controlled trials and observational studies. *Acta psychiatrica Scandinavica*. 2017;135(2):106-116.
43. Vlachos KG, S.; Efremidis, M.; Sideris, A.; Letsas, K. P. An update on risk factors for drug-induced arrhythmias. *Expert review of clinical pharmacology*. 2016;9(1):117-127.
44. Werneke UJ, F.; Taylor, D. M.; Ott, M. Conundrums in neurology: diagnosing serotonin syndrome - a meta-analysis of cases. *BMC neurology*. 2016;16:97.
45. Warden SJF, R. K. Do Selective Serotonin Reuptake Inhibitors (SSRIs) Cause Fractures? *Current osteoporosis reports*. 2016;14(5):211-218.
46. Torjesen I. SSRIs double the risk of suicide and violence in healthy adults. *BMJ (Online)*. 2016;355.
47. Stone M. Suicidality and aggression during antidepressant treatment: Authors misinterpreted earlier paper from the FDA. *BMJ (Online)*. 2016;352.
48. Smit MD, K. M.; Honig, A. Mirtazapine in pregnancy and lactation - A systematic review. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology*. 2016;26(1):126-135.
49. Simonsen ALD, P. B.; Gotzsche, P. C. Persistent sexual dysfunction after early exposure to SSRIs: Systematic review of animal studies. *The International journal of risk & safety in medicine*. 2016;28(1):1-12.
50. Sharma TG, L. S.; Freund, N.; Gotzsche, P. C. Suicidality and aggression during antidepressant treatment: systematic review and meta-analyses based on clinical study reports. *BMJ (Clinical research ed)*. 2016;352:i65.
51. Selmer RH, B.; Furu, K.; Andersen, M.; Norgaard, M.; Zoega, H.; Kieler, H. Individual-based versus aggregate meta-analysis in multi-database studies of pregnancy outcomes: the Nordic example of selective serotonin reuptake inhibitors and venlafaxine in pregnancy. *Pharmacoepidemiology and drug safety*. 2016;25(10):1160-1169.
52. Rudisill TMZ, M.; Kelley, G. A.; Pilkerton, C.; Rudisill, B. R. Medication use and the risk of motor vehicle collisions among licensed drivers: A systematic review. *Accident; analysis and prevention*. 2016;96:255-270.
53. Pozzi MR, S.; Clementi, E.; Molteni, M.; Nobile, M. Antidepressants and, suicide and self-injury: Causal or casual association? *International journal of psychiatry in clinical practice*. 2016;20(1):47-51.
54. Potočnjak ID, V.; Vukičević Baudoin, D.; Čulig, J.; Jakovljević, M. Cardiovascular side effects of psychopharmacologic therapy. *International journal of cardiology*. 2016;219:367-372.
55. Muzik MH, S. E. Use of Antidepressants During Pregnancy?: What to Consider when Weighing Treatment with Antidepressants Against Untreated Depression. *Maternal and child health journal*. 2016;20(11):2268-2279.

56. Lassen D, Ennis ZN, Damkier P. First-Trimester Pregnancy Exposure to Venlafaxine or Duloxetine and Risk of Major Congenital Malformations: A Systematic Review. *Basic & clinical pharmacology & toxicology*. 2016;118(1):32-36.
57. Kaplan YC, Keskin-Arslan E, Acar S, Sozmen K. Prenatal selective serotonin reuptake inhibitor use and the risk of autism spectrum disorder in children: A systematic review and meta-analysis. *Reproductive toxicology (Elmsford, NY)*. 2016;66:31-43.
58. Kobayashi T, Matsuyama T, Takeuchi M, Ito S. Autism spectrum disorder and prenatal exposure to selective serotonin reuptake inhibitors: A systematic review and meta-analysis. *Reproductive toxicology (Elmsford, NY)*. 2016;65:170-178.
59. Jordan SM, J. K.; Davies, G. I.; Tucker, D.; Thayer, D. S.; Luteijn, J. M.; Morgan, M.; Garne, E.; Hansen, A. V.; Klungsoyr, K.; Engeland, A.; Boyle, B.; Dolk, H. Selective Serotonin Reuptake Inhibitor (SSRI) Antidepressants in Pregnancy and Congenital Anomalies: Analysis of Linked Databases in Wales, Norway and Funen, Denmark. *PloS one*. 2016;11(12):e0165122.
60. Jarde AM, M.; Kingston, D.; Giallo, R.; MacQueen, G. M.; Giglia, L.; Beyene, J.; Wang, Y.; McDonald, S. D. Neonatal Outcomes in Women With Untreated Antenatal Depression Compared With Women Without Depression: A Systematic Review and Meta-analysis. *JAMA psychiatry*. 2016;73(8):826-837.
61. Howland RH. Antidepressant Drugs and the Risk of Intracranial Bleeding: Parsing an Observational Study. *Journal of psychosocial nursing and mental health services*. 2016;54(2):21-24.
62. Gallig BR, A.; Nielsen, R. E.; Nielsen, J.; Gerhard, T.; Carbon, M.; Stubbs, B.; Vancampfort, D.; De Hert, M.; Olfson, M.; Kahl, K. G.; Martin, A.; Guo, J. J.; Lane, H. Y.; Sung, F. C.; Liao, C. H.; Arango, C.; Correll, C. U. Type 2 Diabetes Mellitus in Youth Exposed to Antipsychotics: A Systematic Review and Meta-analysis. *JAMA psychiatry*. 2016;73(3):247-259.
63. Donneyong MMB, K.; Bosco-Levy, P.; Dong, Y. H.; Levin, R.; Gagne, J. J. Risk of mortality with concomitant use of tamoxifen and selective serotonin reuptake inhibitors: multi-database cohort study. *BMJ (Clinical research ed)*. 2016;354:i5014.
64. Cameron DHR, M. J. Antidepressants and Driving in Older Adults: A Systematic Review. *Canadian journal on aging = La revue canadienne du vieillissement*. 2016;35 Suppl 1:7-14.
65. Braun CB, T.; Franklin, J.; Baethge, C. Suicides and Suicide Attempts during Long-Term Treatment with Antidepressants: A Meta-Analysis of 29 Placebo-Controlled Studies Including 6,934 Patients with Major Depressive Disorder. *Psychotherapy and psychosomatics*. 2016;85(3):171-179.
66. Bielefeldt AOD, P. B.; Gotzsche, P. C. Precursors to suicidality and violence on antidepressants: systematic review of trials in adult healthy volunteers. *Journal of the Royal Society of Medicine*. 2016;109(10):381-392.
67. Barth MK, L.; Klostermann, S.; Barbui, C.; Cipriani, A.; Linde, K. Efficacy of selective serotonin reuptake inhibitors and adverse events: meta-regression and mediation analysis of placebo-controlled trials. *The British journal of psychiatry : the journal of mental science*. 2016;208(2):114-119.
68. Akioyamen LEM, H.; Holloway, A. C.; Taylor, V. H.; Akioyamen, N. O.; Sherifali, D. Effects of depression pharmacotherapy in fertility treatment on conception, birth, and neonatal health: A systematic review. *Journal of psychosomatic research*. 2016;84:69-80.
69. Alvares GAQ, D. S.; Hickie, I. B.; Guastella, A. J. Autonomic nervous system dysfunction in psychiatric disorders and the impact of psychotropic medications: a systematic review and meta-analysis. *Journal of psychiatry & neuroscience : JPN*. 2016;41(2):89-104.
70. Wang SY, L.; Wang, L.; Gao, L.; Xu, B.; Xiong, Y. Selective Serotonin Reuptake Inhibitors (SSRIs) and the Risk of Congenital Heart Defects: A Meta-Analysis of Prospective Cohort Studies. *Journal of the American Heart Association*. 2015;4(5).
71. Uchida MS, Andrea E.; Biederman, Joseph; Castro, Victor M.; Kenworthy, Tara; Chan, James; Rosales, Ana M.; Newton-Cheh, Christopher; Perlis, Roy H. A systematic evaluation of the QTc interval and antidepressants in youth: An electronic health record study. *Journal of Developmental and Behavioral Pediatrics*. 2015;36(6):434-439.

72. Thase MEF, R.; Cheng, R. F.; Guico-Pabia, C. J.; Sporn, J.; Boucher, M.; Tourian, K. A. Effects of desvenlafaxine on blood pressure in patients treated for major depressive disorder: a pooled analysis. *Current medical research and opinion*. 2015;31(4):809-820.
73. Teo IS, C. T. Assessing the Risks Associated with Antidepressant Use in Plastic Surgery: A Systematic Review. *Plastic and reconstructive surgery*. 2015;136(5):1107-1119.
74. Tampi RRB, M.; Carter, K. V.; Tampi, D. J.; Moca, M.; Knudsen, A.; May, J. Citalopram, QTc Prolongation, and Torsades de Pointes. *Psychosomatics*. 2015;56(1):36-43.
75. Stubbs B. A meta-analysis investigating falls in older adults taking selective serotonin reuptake inhibitors confirms an association but by no means implies causation. *The American Journal of Geriatric Psychiatry*. 2015;23(10):1098-1098.
76. Santarsieri DS, T. L. Antidepressant efficacy and side-effect burden: A quick guide for clinicians. *Drugs in Context*. 2015;4.
77. Robinson GE. Controversies about the use of antidepressants in pregnancy: Response to Dr. Casper, Dr. Osborne, and Dr. Payne. *Journal of Nervous and Mental Disease*. 2015;203(6):480-480.
78. Renoux CL, L. M.; Patenaude, V.; Bresee, L. C.; Paterson, J. M.; Lafrance, J. P.; Tamim, H.; Mahmud, S. M.; Alsabbagh, M. W.; Hemmelgarn, B.; Dormuth, C. R.; Ernst, P. Serotonin-Norepinephrine Reuptake Inhibitors and the Risk of AKI: A Cohort Study of Eight Administrative Databases and Meta-Analysis. *Clinical journal of the American Society of Nephrology : CJASN*. 2015;10(10):1716-1722.
79. Pinheiro EB, Debra L.; Hoxha, Denada; Ciolino, Jody D.; Wisner, Katherine L. Sertraline and breastfeeding: Review and meta-analysis. *Archives of women's mental health*. 2015;18(2):139-146.
80. Orsolini LB, C. Serotonin reuptake inhibitors and breastfeeding: a systematic review. *Human psychopharmacology*. 2015;30(1):4-20.
81. Olfson MG, T. Re-examining antidepressant risk of self-injury in children and adolescents. *Pharmacoepidemiology and drug safety*. 2015;24(2):215-217.
82. Nezafati MHV, M.; Nezafati, P. Antidepressants and cardiovascular adverse events: A narrative review. *ARYA atherosclerosis*. 2015;11(5):295-304.
83. Mohler JF, M.; Chen, Z.; Lee, J. K.; Thienhaus, O. J. Depression, antidepressants, and bone health in older adults: a systematic review. *Journal of the American Geriatrics Society*. 2015;63(3):623-624.
84. Man KKT, H. H.; Wong, L. Y.; Chan, E. W.; Simonoff, E.; Wong, I. C. Exposure to selective serotonin reuptake inhibitors during pregnancy and risk of autism spectrum disorder in children: a systematic review and meta-analysis of observational studies. *Neuroscience and biobehavioral reviews*. 2015;49:82-89.
85. KoKoAung E, Cavenett S, McArthur A, Aromataris E. The association between suicidality and treatment with selective serotonin reuptake inhibitors in older people with major depression: a systematic review. *JB I database of systematic reviews and implementation reports*. 2015;13(3):174-205.
86. Gøtzsche PCY, A. H.; Crace, J. Does long term use of psychiatric drugs cause more harm than good? *BMJ (Online)*. 2015;350.
87. Gentile S. Prenatal antidepressant exposure and the risk of autism spectrum disorders in children. Are we looking at the fall of Gods? *Journal of affective disorders*. 2015;182:132-137.
88. Gentile S. Early pregnancy exposure to selective serotonin reuptake inhibitors, risks of major structural malformations, and hypothesized teratogenic mechanisms. *Expert opinion on drug metabolism & toxicology*. 2015;11(10):1585-1597.
89. Gebara MAL, K. L.; Karp, J. F.; Nash, M. C.; Iaboni, A.; Lenze, E. J. Cause or Effect? Selective Serotonin Reuptake Inhibitors and Falls in Older Adults: A Systematic Review. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2015;23(10):1016-1028.
90. Ennis ZND, P. Pregnancy exposure to olanzapine, quetiapine, risperidone, aripiprazole and risk of congenital malformations. A systematic review. *Basic & clinical pharmacology & toxicology*. 2015;116(4):315-320.
91. Davis JMR, C. E. Do antipsychotics cause hip fractures? Promise and pitfalls of big data. *Journal of Clinical Psychiatry*. 2015;76(9):e1155-e1156.

92. Correll CUD, J.; De Lepeleire, J.; De Hert, M. Effects of antipsychotics, antidepressants and mood stabilizers on risk for physical diseases in people with schizophrenia, depression and bipolar disorder. *World psychiatry : official journal of the World Psychiatric Association (WPA)*. 2015;14(2):119-136.
93. Bruning AHH, H. M.; Kieviet, N.; Bakker, P. C.; de Groot, C. J.; Dolman, K. M.; Honig, A. Antidepressants during pregnancy and postpartum hemorrhage: a systematic review. *European journal of obstetrics, gynecology, and reproductive biology*. 2015;189:38-47.
94. Stevenson JMW, J. L.; Burnham, T. G.; Prevost, A. T.; Schiff, R.; Erskine, S. D.; Davies, J. G. Predicting adverse drug reactions in older adults; A systematic review of the risk prediction models. *Clinical Interventions in Aging*. 2014;9:1581-1593.
95. Ross LEG, Sophie. Selected pregnancy and delivery outcomes after exposure to antidepressant medication. *JAMA psychiatry*. 2014;71(6):716-717.
96. Rais TBR, Alexandra. Association between antidepressants use during pregnancy and autistic spectrum disorders: A meta-analysis. *Innovations in clinical neuroscience*. 2014;11(5-6):18-22.
97. Park HS, H.; Miki, A.; Urushihara, H.; Sawada, Y. Medications associated with falls in older people: systematic review of publications from a recent 5-year period. *European journal of clinical pharmacology*. 2015;71(12):1429-1440.
98. Prieto-Alhambra DP, H.; Goldenberg, J. S.; Khong, T. P.; Klungel, O. H.; Robinson, N. J.; de Vries, F. Excess risk of hip fractures attributable to the use of antidepressants in five European countries and the USA. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2014;25(3):847-855.
99. Previti GP, S.; Chowdhury, S.; Aguglia, E.; Pariante, C. M. Neurodevelopmental outcome for offspring of women treated for antenatal depression: a systematic review. *Archives of women's mental health*. 2014;17(6):471-483.
100. Pereira AC, Y.; Gitlin, M. J.; Dworkin, R. H. Suicidal ideation and behavior associated with antidepressant medications: Implications for the treatment of chronic pain. *Pain*. 2014;155(12):2471-2475.
101. Pedersen LH. Meta-analysis: antidepressant exposure during pregnancy is associated with poor neonatal adaptation. *Evidence-based medicine*. 2014;19(2):76.
102. Paumgartten FJ. Developmental risks associated with use of psychoactive drugs during pregnancy are largely unknown. *Revista Brasileira de Psiquiatria*. 2014;36(4):359.
103. Okazaki R. [Fracture risk associated with drugs other than glucocorticoids]. *Clinical calcium*. 2014;24(3):357-365.
104. Oka YO, K.; Kawashita, N.; Shirakuni, Y.; Takagi, T. Meta-analysis of the risk of upper gastrointestinal hemorrhage with combination therapy of selective serotonin reuptake inhibitors and non-steroidal anti-inflammatory drugs. *Biological & pharmaceutical bulletin*. 2014;37(6):947-953.
105. McDonagh MSM, A.; Phillipi, C.; Romm, J.; Peterson, K.; Thakurta, S.; Guise, J. M. Depression drug treatment outcomes in pregnancy and the postpartum period: a systematic review and meta-analysis. *Obstetrics and gynecology*. 2014;124(3):526-534.
106. Mahdanian AAR, S.; Bacon, S. L.; Ozdin, D.; Lavoie, K. L.; Looper, K. Serotonergic antidepressants and perioperative bleeding risk: a systematic review. *Expert opinion on drug safety*. 2014;13(6):695-704.
107. Mago RT, N.; Andrade, C. Cardiovascular adverse effects of newer antidepressants. *Expert review of neurotherapeutics*. 2014;14(5):539-551.
108. Lorenzo LE, A. Antidepressant use in pregnancy: an evaluation of adverse outcomes excluding malformation. *The Israel journal of psychiatry and related sciences*. 2014;51(2):94-104.
109. Li WL, W.; Wan, Y.; Ren, J.; Li, T.; Li, C. Appraisal of the methodological quality and summary of the findings of systematic reviews on the relationship between SSRIs and suicidality. *Shanghai archives of psychiatry*. 2014;26(5):248-258.
110. Isacsson GR, C. L. Antidepressant drugs and the risk of suicide in children and adolescents. *Paediatric drugs*. 2014;16(2):115-122.

111. Grigoriadis S. The effects of antidepressant medications on mothers and babies. *Journal of population therapeutics and clinical pharmacology = Journal de la therapeutique des populations et de la pharmacologie clinique*. 2014;21(3):e533-541.
112. Gebara MAS, M. L.; Lipsey, K. L.; Teitelbaum, S. L.; Civitelli, R.; Muller, D. J.; Reynolds, C. F., 3rd; Mulsant, B. H.; Lenze, E. J. Depression, antidepressants, and bone health in older adults: a systematic review. *Journal of the American Geriatrics Society*. 2014;62(8):1434-1441.
113. Fanoe SK, D.; Fink-Jensen, A.; Jensen, H. K.; Toft, E.; Nielsen, J.; Videbech, P.; Pehrson, S.; Bundgaard, H. Risk of arrhythmia induced by psychotropic medications: A proposal for clinical management. *European Heart Journal*. 2014;35(20):1306-1315b.
114. El Marroun HW, T.; Verhulst, F. C.; Tiemeier, H. Maternal use of antidepressant or anxiolytic medication during pregnancy and childhood neurodevelopmental outcomes: a systematic review. *European child & adolescent psychiatry*. 2014;23(10):973-992.
115. Costoloni GP, E.; Goracci, A.; Bolognesi, S.; Fagiolini, A. [Mood stabilisers and pregnancy outcomes - a review]. *Psychiatria polska*. 2014;48(5):865-887.
116. Clayton AHC, H. A.; Handiwala, L. Antidepressants and sexual dysfunction: Mechanisms and clinical implications. *Postgraduate medicine*. 2014;126(2):91-99.
117. Beach SRK, W. J.; Celano, C. M.; Januzzi, J. L.; Ruskin, J. N.; Noseworthy, P. A.; Huffman, J. C. Meta-analysis of selective serotonin reuptake inhibitor-associated QTc prolongation. *The Journal of clinical psychiatry*. 2014;75(5):e441-449.
118. Anglin RY, Y.; Moayyedi, P.; Tse, F.; Armstrong, D.; Leontiadis, G. I. Risk of upper gastrointestinal bleeding with selective serotonin reuptake inhibitors with or without concurrent nonsteroidal anti-inflammatory use: a systematic review and meta-analysis. *The American journal of gastroenterology*. 2014;109(6):811-819.
119. Andrade C. Antidepressants and testicular cancer: cause versus association. *The Journal of clinical psychiatry*. 2014;75(3):e198-200.
120. Rotella FM, E. Depression as a risk factor for diabetes: a meta-analysis of longitudinal studies. *The Journal of clinical psychiatry*. 2013;74(1):31-37.
121. Rabenda VN, D.; Beaudart, C.; Bruyere, O.; Reginster, J. Y. Relationship between use of antidepressants and risk of fractures: a meta-analysis. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2013;24(1):121-137.
122. Painuly NP, Ritu; Heun, Reinhard; Sharan, Pratap. Risk of cardiovascular malformations after exposure to paroxetine in pregnancy: Meta-analysis. *The Psychiatrist*. 2013;37(6):198-203.
123. Nassir Ghaemi SV, P. A.; Whitham, E. A. Antidepressants from a public health perspective: Re-examining effectiveness, suicide, and carcinogenicity. *Acta psychiatrica Scandinavica*. 2013;127(2):89-93.
124. Myles NN, H.; Ward, H.; Large, M. Systematic meta-analysis of individual selective serotonin reuptake inhibitor medications and congenital malformations. *The Australian and New Zealand journal of psychiatry*. 2013;47(11):1002-1012.
125. Mintzes B. A leap of faith in antidepressant treatment? *JAMA psychiatry*. 2013;70(12):1-1.
126. Kennedy D. Choosing an SSRI in pregnancy: Clinical context versus statistical significance. *Australian and New Zealand Journal of Psychiatry*. 2013;47(12):1101-1103.
127. Howland RH. Antidepressant medication and pregnancy: time for randomized controlled trials. *J Psychosoc Nurs Ment Health Serv*. 2013;51(2):11-14.
128. Grigoriadis SV, E. H.; Mamisashvili, L.; Roerecke, M.; Rehm, J.; Dennis, C. L.; Koren, G.; Steiner, M.; Mousmanis, P.; Cheung, A.; Ross, L. E. Antidepressant exposure during pregnancy and congenital malformations: is there an association? A systematic review and meta-analysis of the best evidence. *The Journal of clinical psychiatry*. 2013;74(4):e293-308.
129. Grigoriadis SM, L.; Ross, L. E. Achieving the balance: treating depressed pregnant women with antidepressants. *The Journal of clinical psychiatry*. 2013;74(4):375-376.
130. Gahr MS-L, C.; Kölle, M. A.; Freudenmann, R. W. Intoxications with the monoamine oxidase inhibitor tranylcypromine: An analysis of fatal and non-fatal events. *European Neuropsychopharmacology*. 2013;23(11):1364-1372.

131. de Jong MRVDE, M.; Hartholt, K. A. Drug-related falls in older patients: Implicated drugs, consequences, and possible prevention strategies. *Therapeutic Advances in Drug Safety*. 2013;4(4):147-154.
132. de Groot MHvC, J. P.; Moek, M. A.; Tulner, L. R.; Beijnen, J. H.; Lamothe, C. J. The effects of fall-risk-increasing drugs on postural control: a literature review. *Drugs & aging*. 2013;30(11):901-920.
133. Brunnauer AL, G. The effects of most commonly prescribed second generation antidepressants on driving ability: A systematic review: 70th Birthday Prof. Riederer. *Journal of Neural Transmission*. 2013;120(1):225-232.
134. Bhattacharjee SB, R.; Kelley, G. A.; Sambamoorthi, U. Antidepressant use and new-onset diabetes: a systematic review and meta-analysis. *Diabetes/metabolism research and reviews*. 2013;29(4):273-284.
135. Barnard KP, R. C.; Holt, R. I. Antidepressant medication as a risk factor for type 2 diabetes and impaired glucose regulation: systematic review. *Diabetes care*. 2013;36(10):3337-3345.
136. Al-Zoairy RR, C.; Tschoner, A.; Kaser, S.; Ebenbichler, C. The effects of psychotropic drugs on the regulation of glucose metabolism. *Current diabetes reviews*. 2013;9(5):362-370.
137. Wu QB, A. F.; Hentz, J. G.; Crowell, M. D. Selective serotonin reuptake inhibitor treatment and risk of fractures: a meta-analysis of cohort and case-control studies. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2012;23(1):365-375.
138. t Jong GWE, T.; Koren, G.; Einarson, A. Antidepressant use in pregnancy and persistent pulmonary hypertension of the newborn (PPHN): a systematic review. *Reproductive toxicology (Elmsford, NY)*. 2012;34(3):293-297.
139. Sansone RAS, L. A. SSRIs: bad to the bone? *Innovations in clinical neuroscience*. 2012;9(7-8):42-47.
140. Pan AS, Q.; Okereke, O. I.; Rexrode, K. M.; Rubin, R. R.; Lucas, M.; Willett, W. C.; Manson, J. E.; Hu, F. B. Use of antidepressant medication and risk of type 2 diabetes: results from three cohorts of US adults. *Diabetologia*. 2012;55(1):63-72.
141. Oyeboode FR, A.; Berrisford, G.; Coccia, F. Psychotropics in pregnancy: safety and other considerations. *Pharmacology & therapeutics*. 2012;135(1):71-77.
142. Nischal AT, A.; Nischal, A.; Trivedi, J. K. Suicide and antidepressants: What current evidence indicates. *Mens sana monographs*. 2012;10(1):33-44.
143. Malm H. Prenatal exposure to selective serotonin reuptake inhibitors and infant outcome. *Therapeutic drug monitoring*. 2012;34(6):607-614.
144. Lopez-Yarto MR-M, E.; Holloway, A. C.; Taylor, V. H.; McDonald, S. D. Do psychiatric medications, especially antidepressants, adversely impact maternal metabolic outcomes? *Journal of affective disorders*. 2012;141(2-3):120-129.
145. Kucukaycan MVDE, F.; Moreels, T.; Sabbe, B. G. [Antidepressants and the risk of bleeding]. *Tijdschrift voor psychiatrie*. 2012;54(3):225-234.
146. KoKoAung EA, E. The association between suicidality and treatment with selective serotonin reuptake inhibitors in older people with major depression: A systematic review. *JBIC library of systematic reviews*. 2012;10(56 SUPPL):S313-S324.
147. Hennings JMS, L.; Fulda, S. Glucose metabolism and antidepressant medication. *Current pharmaceutical design*. 2012;18(32):5900-5919.
148. Hackam DGM, M. Selective serotonin reuptake inhibitors and brain hemorrhage: a meta-analysis. *Neurology*. 2012;79(18):1862-1865.
149. Grzeskowiak LEG, A. L.; Morrison, J. L. Exposed or not exposed? Exploring exposure classification in studies using administrative data to investigate outcomes following medication use during pregnancy. *European journal of clinical pharmacology*. 2012;68(5):459-467.
150. Einarson A. Publishing statistically significant results with questionable clinical importance: focus on antidepressant use in pregnancy. *The Journal of clinical psychiatry*. 2012;73(11):1443-1446.
151. Eom CS, Lee HK, Ye S, Park SM, Cho KH. Use of selective serotonin reuptake inhibitors and risk of fracture: a systematic review and meta-analysis. *J Bone Miner Res*. 2012;27(5):1186-1195.

152. Colotto MV, F.; Vo Hong, N.; Raimo, O.; Castello, A.; Carnovale, A.; Paciaroni, A.; Coletta, P. [Effect of treatment with selective serotonin reuptake inhibitors on lipid profile: state of the art]. *La Clinica terapeutica*. 2012;163(1):e41-45.
153. Bromley RLW, A.; Makarova, D.; Tower, C.; Wood, A.; Clayton-Smith, J. Fetal effects of selective serotonin reuptake inhibitor treatment during pregnancy: Immediate and longer term child outcomes. *Fetal and Maternal Medicine Review*. 2012;23(3-4):230-275.
154. Adams KB, C.; van der Heijde, D. M. Safety of pain therapy during pregnancy and lactation in patients with inflammatory arthritis: a systematic literature review. *The Journal of rheumatology Supplement*. 2012;90:59-61.
155. Gentile SG, M. Prenatal exposure to antidepressant medications and neurodevelopmental outcomes: a systematic review. *Journal of affective disorders*. 2011;128(1-2):1-9.
156. Gentile S. Neonatal withdrawal reactions following late in utero exposure to antidepressant medications. *Current Women's Health Reviews*. 2011;7(1):18-27.
157. Fenger-Gron JT, M.; Andersen, K. S.; Nielsen, R. G. Paediatric outcomes following intrauterine exposure to serotonin reuptake inhibitors: a systematic review. *Danish medical bulletin*. 2011;58(9):A4303.
158. Davanzo RC, M.; De Cunto, A.; Minen, F.; Amaddeo, A. Antidepressant drugs and breastfeeding: a review of the literature. *Breastfeeding medicine : the official journal of the Academy of Breastfeeding Medicine*. 2011;6(2):89-98.
159. Dassanayake TM, P.; Carter, G.; Jones, A. Effects of benzodiazepines, antidepressants and opioids on driving: a systematic review and meta-analysis of epidemiological and experimental evidence. *Drug safety*. 2011;34(2):125-156.
160. Berkowitz RIF, A. N. Obesity, psychiatric status, and psychiatric medications. *Psychiatric Clinics of North America*. 2011;34(4):747-764.
161. Barbui C. Getting through the quicksand of the relationship between drugs and suicide. *Drug safety*. 2011;34(5):397-401.
162. Wurst KEP, C.; Ephross, S. A.; Olshan, A. F. First trimester paroxetine use and the prevalence of congenital, specifically cardiac, defects: a meta-analysis of epidemiological studies. *Birth defects research Part A, Clinical and molecular teratology*. 2010;88(3):159-170.
163. Wu QL, J.; Gallegos-Orozco, J. F.; Hentz, J. G. Depression, fracture risk, and bone loss: a meta-analysis of cohort studies. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2010;21(10):1627-1635.
164. van Driel JJW, J. H.; Heres, M. H.; Honig, A. [Intrauterine exposure to SSRIs: no reason for routine hypoglycaemia screening]. *Nederlands tijdschrift voor geneeskunde*. 2010;154(8):A339.
165. Udechuku AN, T.; Hill, R.; Szego, K. Antidepressants in pregnancy: a systematic review. *The Australian and New Zealand journal of psychiatry*. 2010;44(11):978-996.
166. Tuccori MM, S.; Testi, A.; Ruggiero, E.; Mantarro, S.; Scollo, C.; Pergola, A.; Fornai, M.; Antonioli, L.; Colucci, R.; Corona, T.; Blandizzi, C. Use of selective serotonin reuptake inhibitors during pregnancy and risk of major and cardiovascular malformations: an update. *Postgraduate medicine*. 2010;122(4):49-65.
167. Simoncelli MM, B. Z.; Berard, A. Antidepressant use during pregnancy: a critical systematic review of the literature. *Current drug safety*. 2010;5(2):153-170.
168. Scialli AR. Paroxetine exposure during pregnancy and cardiac malformations. *Birth defects research Part A, Clinical and molecular teratology*. 2010;88(3):175-177.
169. Kemp AH, Quintana DS, Gray MA, Felmingham KL, Brown K, Gatt JM. Impact of depression and antidepressant treatment on heart rate variability: a review and meta-analysis. *Biological psychiatry*. 2010;67(11):1067-1074.
170. Kölch MP, P.; Allroggen, M.; Fegert, J. M. Does behavioural toxicity of SSRI increased risk for school shootings? *Nervenheilkunde*. 2010;29(7-8):451-455.
171. Kontakioti EB, Z.; Tsiaousi, M.; Mironidou-Tzouveleki, M. Selective serotonin reuptake inhibitors in pregnancy and breastfeeding. *Epitheorese Klinikes Farmakologias kai Farmakokinetikes*. 2010;28(3):249-268.

172. Einarson AK, G. First trimester exposure to paroxetine and prevalence of cardiac defects: Meta-analysis of the literature: Unfortunately incomplete. *Birth defects research Part A, Clinical and molecular teratology*. 2010;88(7):588; author reply 589.
173. Einarson A. Paroxetine use in pregnancy and increased risk of heart defects: Evaluating the evidence. *Canadian family physician Medecin de famille canadien*. 2010;56(8):767-768.
174. Berard A. Paroxetine exposure during pregnancy and the risk of cardiac malformations: what is the evidence? *Birth defects research Part A, Clinical and molecular teratology*. 2010;88(3):171-174.
175. Woolcott JCR, K. J.; Wiens, M. O.; Patel, B.; Marin, J.; Khan, K. M.; Marra, C. A. Meta-analysis of the impact of 9 medication classes on falls in elderly persons. *Archives of internal medicine*. 2009;169(21):1952-1960.
176. Fortinguerra FC, A.; Bonati, M. Psychotropic drug use during breastfeeding: A review of the evidence. *Pediatrics*. 2009;124(4):e547-e556.
177. Bergemann N. Antidepressant use during pregnancy. *Journal fur Neurologie, Neurochirurgie und Psychiatrie*. 2009;10(1):41-47.
178. Taylor D. Antidepressant drugs and cardiovascular pathology: A clinical overview of effectiveness and safety. *Acta psychiatrica Scandinavica*. 2008;118(6):434-442.
179. Sterke CSV, A. P.; van Beeck, E. F.; van der Cammen, T. J. The influence of drug use on fall incidents among nursing home residents: a systematic review. *International psychogeriatrics*. 2008;20(5):890-910.
180. O'Brien LE, T. R.; Sarkar, M.; Einarson, A.; Koren, G. Does paroxetine cause cardiac malformations? *Journal of obstetrics and gynaecology Canada : JOGC = Journal d'obstetrique et gynecologie du Canada : JOGC*. 2008;30(8):696-701.
181. Loke YKT, A. N.; Singh, S. Meta-analysis: gastrointestinal bleeding due to interaction between selective serotonin uptake inhibitors and non-steroidal anti-inflammatory drugs. *Alimentary pharmacology & therapeutics*. 2008;27(1):31-40.
182. Kongkaew CN, P. R.; Ashcroft, D. M. Hospital admissions associated with adverse drug reactions: A systematic review of prospective observational studies. *Annals of Pharmacotherapy*. 2008;42(7-8):1017-1025.
183. Bond DJN, M. M.; Kauer-Sant'Anna, M.; Lam, R. W.; Yatham, L. N. Antidepressant-associated mood elevations in bipolar II disorder compared with bipolar I disorder and major depressive disorder: a systematic review and meta-analysis. *The Journal of clinical psychiatry*. 2008;69(10):1589-1601.
184. Takkouche BM-M, A.; Gill, S. S.; Etminan, M. Psychotropic medications and the risk of fracture: a meta-analysis. *Drug safety*. 2007;30(2):171-184.
185. Howard RL, Avery AJ, Slavenburg S, et al. Which drugs cause preventable admissions to hospital? A systematic review. *British journal of clinical pharmacology*. 2007;63(2):136-147.
186. Hartikainen SL, E.; Louhivuori, K. Medication as a risk factor for falls: critical systematic review. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2007;62(10):1172-1181.
187. Bellantuono CM, G.; Gentile, S. Serotonin reuptake inhibitors in pregnancy and the risk of major malformations: a systematic review. *Human psychopharmacology*. 2007;22(3):121-128.
188. Bar-Oz BE, T.; Einarson, A.; Boskovic, R.; O'Brien, L.; Malm, H.; Berard, A.; Koren, G. Paroxetine and congenital malformations: meta-Analysis and consideration of potential confounding factors. *Clinical therapeutics*. 2007;29(5):918-926.
189. Swenson JRD, S.; Fergusson, D. Adverse cardiovascular events in antidepressant trials involving high-risk patients: a systematic review of randomized trials. *Canadian journal of psychiatry Revue canadienne de psychiatrie*. 2006;51(14):923-929.
190. Sala MC, F.; Cappucciati, C.; Brambilla, P.; d'Allio, G.; Caverzasi, E.; Barale, F.; De Ferrari, G. Antidepressants: Their effects on cardiac channels, QT prolongation and Torsade de Pointes. *Current Opinion in Investigational Drugs*. 2006;7(3):256-263.
191. Rahimi RN, S.; Abdollahi, M. Pregnancy outcomes following exposure to serotonin reuptake inhibitors: a meta-analysis of clinical trials. *Reproductive toxicology (Elmsford, NY)*. 2006;22(4):571-575.
192. McIntyre RSS, J. K.; Konarski, J. Z.; Kennedy, S. H. The effect of antidepressants on lipid homeostasis: a cardiac safety concern? *Expert opinion on drug safety*. 2006;5(4):523-537.

193. McClintock SMH, M. M. Association of suicide in the elderly and SSRI antidepressants: Information from a population-based study. *Aging Health*. 2006;2(6):917-918.
194. Hall WDL, J. How have the selective serotonin reuptake inhibitor antidepressants affected suicide mortality? *The Australian and New Zealand journal of psychiatry*. 2006;40(11-12):941-950.
195. Lattimore KAD, S. M.; Kaciroti, N.; Kemper, A. R.; Neal, C. R., Jr.; Vazquez, D. M. Selective serotonin reuptake inhibitor (SSRI) use during pregnancy and effects on the fetus and newborn: a meta-analysis. *Journal of perinatology : official journal of the California Perinatal Association*. 2005;25(9):595-604.
196. Gentile S. The safety of newer antidepressants in pregnancy and breastfeeding. *Drug safety*. 2005;28(2):137-152.
197. Einarson TRE, A. Newer antidepressants in pregnancy and rates of major malformations: a meta-analysis of prospective comparative studies. *Pharmacoepidemiology and drug safety*. 2005;14(12):823-827.
198. Wen SWW, M. Risk of fetal exposure to tricyclic antidepressants. *Journal of obstetrics and gynaecology Canada : JOGC = Journal d'obstetrique et gynecologie du Canada : JOGC*. 2004;26(10):887-892.
199. Weissman AML, B. T.; Hartz, A. J.; Bentler, S.; Donohue, M.; Ellingrod, V. L.; Wisner, K. L. Pooled analysis of antidepressant levels in lactating mothers, breast milk, and nursing infants. *The American journal of psychiatry*. 2004;161(6):1066-1078.
200. Ramasubbu R. Cerebrovascular effects of selective serotonin reuptake inhibitors: a systematic review. *The Journal of clinical psychiatry*. 2004;65(12):1642-1653.
201. Newman TB. A black-box warning for antidepressants in children? *The New England journal of medicine*. 2004;351(16):1595-1598.
202. McIntosh HMW, N. F.; Bagnall, A. M. Assessing harmful effects in systematic reviews. *BMC medical research methodology*. 2004;4:19.
203. Bailey BB, N. A.; Amre, D. K. A meta-analysis of prognostic indicators to predict seizures, arrhythmias or death after tricyclic antidepressant overdose. *Journal of toxicology Clinical toxicology*. 2004;42(6):877-888.
204. Weinrieb RMA, Marc; Lynch, Kevin G.; Chang, Kyong-Mi; Lewis, James D. A critical review of selective serotonin reuptake inhibitor-associated bleeding: Balancing the risk of treating Hepatitis C-infected patients. *The Journal of clinical psychiatry*. 2003;64(12):1502-1510.
205. Addis AK, G. Safety of fluoxetine during the first trimester of pregnancy: a meta-analytical review of epidemiological studies. *Psychological medicine*. 2000;30(1):89-94.
206. Thase ME. Effects of venlafaxine on blood pressure: a meta-analysis of original data from 3744 depressed patients. *The Journal of clinical psychiatry*. 1998;59(10):502-508.
207. Wilens TEB, Joseph; Baldessarini, Ross J.; Geller, Barbara; Schleifer, David; Spencer, Thomas J.; Birmaher, Boris; Goldblatt, Alan. Cardiovascular effects of therapeutic doses of tricyclic antidepressants in children and adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1996;35(11):1491-1501.
208. Radua J, Ramella-Cravaro V, Ioannidis JPA, et al. What causes psychosis? An umbrella review of risk and protective factors. *World Psychiatry*. 2018;17(1):49-66.
209. Theodoratou E, Tzoulaki I, Zgaga L, Ioannidis JP. Vitamin D and multiple health outcomes: umbrella review of systematic reviews and meta-analyses of observational studies and randomised trials. *BMJ*. 2014;348:g2035.
210. Raglan O, Kalliala I, Markozannes G, et al. Risk factors for endometrial cancer: An umbrella review of the literature. *Int J Cancer*. 2018.
211. Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017;358:j4008.
212. Perry R, Leach V, Davies P, Penfold C, Ness A. A comparison of two assessment tools used in overviews of systematic reviews: ROBIS versus AMSTAR. In: *Cochrane Colloquium Abstracts 2018*

Edinburgh. Available at <https://abstracts.cochrane.org/comparison-two-assessment-tools-used-overviews-systematic-reviews-robis-versus-amstar>.

213. Ioannidis JPA. Integration of evidence from multiple meta-analyses: a primer on umbrella reviews, treatment networks and multiple treatments meta-analyses. CMAJ. 2009;181(8):488-93. doi: 10.1503/cmaj.08108.
214. Atkins D, Eccles M, Flottorp S, Guyatt GH, Henry D, Hill S, Liberati A, O'Connell D, Oxman AD, Phillips B, Schünemann H, Edejer TT, Vist GE, Williams JW Jr; GRADE Working Group. Systems for grading the quality of evidence and the strength of recommendations I: critical appraisal of existing approaches The GRADE Working Group. BMC Health Serv Res. 2004;4(1):38.
215. Morales DR, Slattery J, Evans S, Kurz X. Antidepressant use during pregnancy and risk of autism spectrum disorder and attention deficit hyperactivity disorder: systematic review of observational studies and methodological considerations. BMC Med. 2018 Jan 15;16(1):6.




Figure 1 Flowchart outlining the literature search and evaluation process of published meta-analyses and systematic reviews




Figure 2. Percentages of the reported 13 adverse health outcomes related to exposure to antidepressants in the published 45 meta-analyses.